

UNIQUE CONTINUATION FOR DISCRETE NONLINEAR WAVE EQUATIONS

HELGE KRÜGER AND GERALD TESCHL

ABSTRACT. We establish unique continuation for various discrete nonlinear wave equations. For example, we show that if two solutions of the Toda lattice coincide for one lattice point in some arbitrarily small time interval, then they coincide everywhere. Moreover, we establish analogous results for the Toda, Kac–van Moerbeke, and Ablowitz–Ladik hierarchies. Although all these equations are integrable, the proof does not use integrability and can be adapted to other equations as well.

1. INTRODUCTION

Unique continuation results for wave equations have a long tradition and seem to originate in control theory. One of the first result seems to be the one by Zhang [20], where he proves that if a short range solution of the Korteweg–de Vries (KdV) equation vanishes on an open subset in the xt -plane, then it must vanish everywhere. Since then, this result has been extended in various directions and for different equations (see for example [2], the introduction in [10] for the case of the nonlinear Schrödinger equation, [5], [11], [12] for the generalized KdV equation, [14] for the Camassa–Holm equation).

However, all the results so far seem to only deal with wave equations which are continuous in the spatial direction and this clearly raises the question for such unique continuation results for wave equations which are discrete in the spatial variable. In particular, to the best of our knowledge there are no results for example for the Toda equation, one of the most prominent discrete system. While in principle the strategy from Zhang [20] would be applicable to the Toda lattice, it is the purpose of this note to advocate a much simpler direct approach in the discrete case. We will start with the Toda lattice as our prototypical example and then show how the entire Toda hierarchy as well as the Kac–van Moerbeke and Ablowitz–Ladik hierarchies can be treated. It is important to stress that our approach does not use integrability of these equations and hence can be adapted to more general systems. On the other hand, our approach seems to need one dimensionality in the spatial variable and thus does not apply to the discrete Schrödinger equation on \mathbb{Z}^d . Due to the connections with localization for discrete Anderson–Bernoulli models, unique continuation for this model is an important open problem; see [3], [4].

2000 *Mathematics Subject Classification.* Primary 35L05, 37K60; Secondary 37K15, 37K10.

Key words and phrases. Unique continuation, Toda lattice, Kac–van Moerbeke lattice, Ablowitz–Ladik equations, discrete nonlinear Schrödinger equation, Schur flow.

Research supported by the Austrian Science Fund (FWF) under Grant No. Y330 and the National Science Foundation (NSF) under Grant No. DMS-0800100.

2. THE TODA LATTICE

In this section we want to treat the Toda lattice as the prototypical example. To this end, recall the Toda lattice [19] (in Flaschka's variables [7])

$$(2.1) \quad \begin{aligned} \dot{a}(n, t) &= a(n, t) \left(b(n+1, t) - b(n, t) \right), \\ \dot{b}(n, t) &= 2 \left(a(n, t)^2 - a(n-1, t)^2 \right), \quad n \in \mathbb{Z}, \end{aligned}$$

where the dot denotes a derivative with respect to t . It is a well studied physical model and one of the prototypical discrete integrable wave equations. We refer to the monographs [6], [16], [19] or the review articles [13], [17] for further information.

Theorem 2.1. *Assume that $a_0(n, t) \neq 0$, $b_0(n, t)$ and $a(n, t), b(n, t)$ are complex-valued solutions of (2.1) such that there is one $n_0 \in \mathbb{Z}$ and two times $t_0 < t_1$ such that*

$$(2.2) \quad a_0(n_0, t)^2 = a(n_0, t)^2, \quad b_0(n_0, t) = b(n_0, t),$$

for $t \in (t_0, t_1)$. Then

$$(2.3) \quad a_0(n, t)^2 = a(n, t)^2, \quad b_0(n, t) = b(n, t)$$

for all $(n, t) \in \mathbb{Z} \times \mathbb{R}$.

Proof. It suffices to prove that (2.2) for n_0 implies (2.2) for $n_0 - 1$ and $n_0 + 1$. We start with $N_0 - 1$ and first observe that (2.1) implies

$$\begin{aligned} 0 &= \dot{b}(n_0, t) - \dot{b}_0(n_0, t) = 2(a(n_0, t)^2 - a_0(n_0, t)^2 - a(n_0 - 1, t)^2 + a_0(n_0 - 1, t)^2) \\ &= -2(a(n_0 - 1, t)^2 - a_0(n_0 - 1, t)^2) \end{aligned}$$

and thus $a(n_0 - 1, t)^2 = a_0(n_0 - 1, t)^2$. Using this we compute

$$\begin{aligned} 0 &= \frac{\dot{a}(n_0 - 1, t)}{a(n_0 - 1, t)} - \frac{\dot{a}_0(n_0 - 1, t)}{a_0(n_0 - 1, t)} \\ &= b(n_0, t) - b_0(n_0, t) - b(n_0 - 1, t) + b_0(n_0 - 1, t) \\ &= -b(n_0 - 1, t) + b_0(n_0 - 1, t), \end{aligned}$$

so $b(n_0 - 1, t) = b_0(n_0 - 1, t)$. Now for $n_0 + 1$, we begin with

$$\begin{aligned} 0 &= \frac{\dot{a}(n_0, t)}{a(n_0, t)} - \frac{\dot{a}_0(n_0, t)}{a_0(n_0, t)} = b(n_0 + 1, t) - b_0(n_0 + 1, t) - b(n_0, t) + b_0(n_0, t) \\ &= b(n_0 + 1, t) - b_0(n_0 + 1, t), \end{aligned}$$

so $b(n_0 + 1, t) = b_0(n_0 + 1, t)$. Now, use that

$$\begin{aligned} 0 &= \dot{b}(n_0 + 1, t) - \dot{b}_0(n_0 + 1, t) \\ &= 2(a(n_0 + 1, t)^2 - a_0(n_0 + 1, t)^2 - a(n_0, t)^2 + a_0(n_0, t)^2) \\ &= 2(a(n_0 + 1, t)^2 - a_0(n_0 + 1, t)^2) \end{aligned}$$

to conclude $a(n_0 + 1, t)^2 = a_0(n_0 + 1, t)^2$. This finishes the proof. \square

It is worthwhile to note that the assumption $a(n, t)^2 \neq 0$ is crucial. In fact, if $a(n_0, t) = 0$ for one (and hence for all) $t \in \mathbb{R}$, then the Toda lattice decouples into two independent parts to the left and right of n_0 , and the above result is clearly wrong.

As a simple consequence, this also proves finite propagation speed for the Toda lattice.

Corollary 2.2. *Let $a(n, t)$, $b(n, t)$ be a solution of the Toda lattice for which $a(n, t_0) - \frac{1}{2}$, $b(n, t_0)$ is compactly supported. Then this does not remain true for $t \in (t_0, t_1)$ unless $a(n, t) = \frac{1}{2}$, $b(n, t) = 0$.*

In fact, in the case of real-valued solutions, one can even show the somewhat stronger result, that $a(n, t_0) - \frac{1}{2}$, $b(n, t_0)$ can be compactly supported for at most one time [18]. However, on the other hand the Toda lattice does preserve certain asymptotic properties of the initial conditions, see again [18].

3. EXTENSION TO THE TODA AND KAC-VAN MOERBEKE HIERARCHY

In this section we show that our main result extends to the entire Toda hierarchy (which will cover the Kac-van Moerbeke hierarchy as well). To this end, we introduce the Toda hierarchy using the standard Lax formalism following [1] (see also [9], [16]).

Associated with two sequences $a(t)^2 \neq 0$, $b(t)$ is a Jacobi operator

$$(3.1) \quad H(t) = a(t)S^+ + a^-(t)S^- + b(t)$$

acting on sequences over \mathbb{Z} , where $S^\pm f(n) = f^\pm(n) = f(n \pm 1)$ are the usual shift operators. Moreover, choose constants $c_0 = 1$, c_j , $1 \leq j \leq r$, $c_{r+1} = 0$, and set

$$(3.2) \quad P_{2r+2}(t) = \sum_{j=0}^r c_{r-j} \tilde{P}_{2j+2}(t), \quad \tilde{P}_{2j+2}(t) = [H(t)^{j+1}]_+ - [H(t)^{j+1}]_-,$$

where $[A]_\pm$ denote the upper and lower triangular parts of an operator with respect to the standard basis $\delta_m(n) = \delta_{m,n}$ (with $\delta_{m,n}$ the usual Kronecker delta). Then the Toda hierarchy is equivalent to the Lax equation

$$(3.3) \quad \frac{d}{dt} H(t) - [P_{2r+2}(t), H(t)] = 0, \quad t \in \mathbb{R},$$

where $[A, B] = AB - BA$ is the usual commutator. Abbreviating

$$(3.4) \quad g_j(n, t) = \sum_{\ell=0}^j c_{j-\ell} \tilde{g}_\ell(n, t), \quad \tilde{g}_\ell(n, t) = \langle \delta_n, H(t)^\ell \delta_n \rangle,$$

$$h_j(n, t) = 2a(n, t) \sum_{\ell=0}^j c_{j-\ell} \tilde{h}_\ell(n, t) + c_{j+1}, \quad \tilde{h}_\ell(n, t) = \langle \delta_{n+1}, H(t)^\ell \delta_n \rangle.$$

one explicitly obtains

$$(3.5) \quad \text{TL}_r(a(t), b(t)) = \begin{pmatrix} \dot{a}(t) - a(t) \left(g_{r+1}^+(t) - g_{r+1}(t) \right) \\ \dot{b}(t) - \left(h_{r+1}(t) - h_{r+1}^-(t) \right) \end{pmatrix} = 0, \quad r \in \mathbb{N}_0,$$

for the r 'th equation $\text{TL}_r(a, b) = 0$ in the Toda hierarchy (where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$).

Our main point in this section is the following generalization of Theorem 2.1 to the entire Toda hierarchy:

Theorem 3.1. *Assume that $a_0(n, t)^2 \neq 0$, $b_0(n, t)$ and $a(n, t)^2, b(n, t)$ are complex-valued solutions of some equation in the Toda hierarchy TL_r such that there is one $n_0 \in \mathbb{Z}$ and two times $t_0 < t_1$ such that*

$$(3.6) \quad a_0(n_0 + j, t)^2 = a(n_0 + j, t)^2, \quad b_0(n_0 + j, t) = b(n_0 + j, t), \quad j = 0, \dots, r,$$

for $t \in (t_0, t_1)$. Then

$$(3.7) \quad a_0(n, t)^2 = a(n, t)^2, \quad b_0(n, t) = b(n, t)$$

for all $(n, t) \in \mathbb{Z} \times \mathbb{R}$.

Proof. Let us drop the dependence on t for notational simplicity during this proof. The key observation is the following structure for the homogenous quantities \tilde{g}_j, \tilde{h}_j :

$$\tilde{g}_j(n) = \begin{cases} \left(\prod_{\ell=0}^{k-1} a(n+\ell)^2 \right) b(n+k) + R(n+k-1, n-k+1) + \\ + \left(\prod_{\ell=1}^k a(n-\ell)^2 \right) \left(b(n-k) + 2 \sum_{\ell=0}^{k-1} b(n-\ell) \right), & j = 2k+1, \\ \left(\prod_{\ell=0}^{k-2} a(n+\ell)^2 \right) \left(a(n+k-1)^2 + b(n+k-1)^2 \right. \\ \left. + 2b(n+k-1) \sum_{\ell=0}^{k-2} b(n+\ell) \right) + \\ + R(n+k-2, n-k+1) + \prod_{\ell=1}^k a(n-\ell)^2, & j = 2k, \end{cases}$$

and

$$\tilde{h}_j(n) = \begin{cases} 2 \left(\prod_{\ell=0}^{k-1} a(n+\ell)^2 \right) \left(a(n+k)^2 + b(n+k)^2 \right. \\ \left. + 2b(n+k) \sum_{\ell=0}^{k-1} b(n+\ell) \right) + \\ + R(n+k-1, n-k+1) + 2 \prod_{\ell=0}^k a(n-\ell)^2, & j = 2k+1, \\ 2 \left(\prod_{\ell=0}^{k-1} a(n+\ell)^2 \right) b(n+k) + R(n+k-1, n-k+2) \\ + 2 \prod_{\ell=0}^{k-1} a(n-\ell)^2 \left(b(n+1) + b(n-k+1) + 2 \sum_{\ell=0}^{k-2} b(n-\ell) \right), & j = 2k, \end{cases}$$

for $j > 1$. Here $R(n, m)$ denotes terms which involve only $a(\ell)$ and $b(\ell)$ with $m \leq \ell \leq n$ and we set $R(n, m) = 0$ if $n < m$. In fact, this can be verified using $\tilde{g}_0 = 1, \tilde{h}_0 = 0$ together with the recursions ([16, Chap. 6])

$$(3.8) \quad \tilde{g}_{j+1} = \frac{\tilde{h}_j + \tilde{h}_j^-}{2} + b\tilde{g}_j,$$

$$(3.9) \quad \tilde{h}_{j+1} = 2a^2 \sum_{\ell=0}^j \tilde{g}_{j-\ell} \tilde{g}_\ell^+ - \frac{1}{2} \sum_{\ell=0}^j \tilde{h}_{j-\ell} \tilde{h}_\ell, \quad j \in \mathbb{N}_0.$$

Now we are ready for the main part of the proof. It suffices to show that (3.7) hold for $n = n_0 - 1$ and $n = n_0 + r + 1$.

We first look at the case $r + 1 = 2k + 1$. Then

$$\begin{aligned} 0 &= \frac{\dot{a}(n_0 + k)}{a(n_0 + k)} - \frac{\dot{a}_0(n_0 + k)}{a_0(n_0 + k)} \\ &= g_{r+1}(n_0 + k + 1) - g_{r+1}(n_0 + k) - g_{0,r+1}(n_0 + k + 1) + g_{0,r+1}(n_0 + k) \\ &= \left(\prod_{\ell=k+1}^r a_0(n_0 + \ell)^2 \right) (b(n_0 + r + 1) - b_0(n_0 + r + 1)) \end{aligned}$$

shows $b(n_0 + r + 1) = b_0(n_0 + r + 1)$. Similarly,

$$\begin{aligned} 0 &= \dot{b}(n_0 + k) - \dot{b}_0(n_0 + k) \\ &= h_{r+1}(n_0 + k) - h_{r+1}(n_0 + k - 1) - h_{0,r+1}(n_0 + k) + h_{0,r+1}(n_0 + k - 1) \\ &= 2 \left(\prod_{\ell=0}^{k-1} a_0(n_0 + \ell)^2 \right) (a(n_0 - 1)^2 - a_0(n_0 - 1)^2) \end{aligned}$$

shows $a(n_0 - 1)^2 = a_0(n_0 - 1)^2$. Proceeding like this, and using the result found in the previous steps,

$$\begin{aligned} 0 &= \dot{b}(n_0 + k + 1) - \dot{b}_0(n_0 + k + 1) \\ &= h_{r+1}(n_0 + k + 1) - h_{r+1}(n_0 + k) - h_{0,r+1}(n_0 + k + 1) + h_{0,r+1}(n_0 + k) \\ &= \left(\prod_{\ell=k+1}^r a_0(n_0 + \ell)^2 \right) (a(n_0 + r + 1)^2 - a_0(n_0 + r + 1)^2) \end{aligned}$$

shows $a(n_0 + r + 1)^2 = a_0(n_0 + r + 1)^2$ and

$$\begin{aligned} 0 &= \frac{\dot{a}(n_0 + k - 1)}{a(n_0 + k - 1)} - \frac{\dot{a}_0(n_0 + k - 1)}{a_0(n_0 + k - 1)} \\ &= g_{r+1}(n_0 + k) - g_{r+1}(n_0 + k - 1) - g_{0,r+1}(n_0 + k) + g_{0,r+1}(n_0 + k - 1) \\ &= \left(\prod_{\ell=-1}^k a_0(n_0 + \ell)^2 \right) (b(n_0 - 1) - b_0(n_0 - 1)) \end{aligned}$$

shows $b(n_0 - 1) = b_0(n_0 - 1)$, which finishes the case $r + 1 = 2k + 1$. The case $r + 1 = 2k$ is analogous. \square

Finally since the Kac-van Moerbeke hierarchy can be obtained by setting $b = 0$ in the odd equations of the Toda hierarchy, $\text{KM}_r(a) = \text{TL}_{2r+1}(a, 0)$ (see [15]), this last result also covers the Kac-van Moerbeke hierarchy. In particular,

Corollary 3.2. *Assume that $\rho_0(n, t) > 0$ and $\rho(n, t) > 0$ are solutions of the Kac-van Moerbeke equation*

$$(3.10) \quad \dot{\rho}(n, t) = \rho(n, t)(\rho(n + 1, t) - \rho(n - 1, t))$$

such that there is one $n_0 \in \mathbb{Z}$ and two times $t_0 < t_1$ such that

$$(3.11) \quad \rho_0(n_0, t) = \rho(n_0, t), \quad \rho_0(n_0 + 1, t) = \rho(n_0 + 1, t),$$

for $t \in (t_0, t_1)$. Then

$$(3.12) \quad \rho_0(n, t) = \rho(n, t)$$

for all $(n, t) \in \mathbb{Z} \times \mathbb{R}$.

4. THE ABLOWITZ–LADIK HIERARCHY

In this section we show that our main result extends to the Ablowitz–Ladik (AL) hierarchy [9]. We first state the result for the simplest case, whose proof follows as the one of Theorem 2.1.

Theorem 4.1. *Let $C_{0,\pm}, c_1 \in \mathbb{C} \setminus \{0\}$. Assume that $\alpha_0(n, t), \beta_0(n, t)$, with $\rho_0(n, t) \neq 0$, and $\alpha(n, t), \beta(n, t)$ are solutions of the Ablowitz–Ladik equation*

$$(4.1) \quad \begin{aligned} i\dot{\alpha}(n, t) &= -\rho(n, t)^2 (c_{0,-}\alpha(n-1, t) + c_{0,+}\alpha(n+1, t)) - c_1\alpha(n, t), \\ i\dot{\beta}(n, t) &= \rho(n, t)^2 (c_{0,+}\beta(n-1, t) + c_{0,-}\beta(n+1, t)) + c_1\beta(n, t), \end{aligned}$$

where

$$(4.2) \quad \rho(n, t) = (1 - \alpha(n, t)\beta(n, t))^{1/2},$$

such that there is one $n_0 \in \mathbb{Z}$ and two times $t_0 < t_1$ such that

$$(4.3) \quad \alpha_0(n_0 + j, t) = \alpha(n_0 + j, t), \quad \beta_0(n_0 + j, t) = \beta(n_0 + j, t), \quad j = 0, 1,$$

for $t \in (t_0, t_1)$. Then

$$(4.4) \quad \alpha_0(n, t) = \alpha(n, t), \quad \beta_0(n, t) = \beta(n, t)$$

for all $(n, t) \in \mathbb{Z} \times \mathbb{R}$.

The special choices $c_{0,\pm} = 1$, $c_1 = -2$, and $\beta = \pm\bar{\alpha}$ yield the focusing, defocusing discrete nonlinear Schrödinger equations, respectively. The alternative choice $c_{0,\pm} = \pm i$, $c_1 = 0$, and $\beta = \bar{\alpha}$ yield the Schur flow.

We next turn to the AL hierarchy following [8], [9]. Associated with two sequences $\alpha(t), \beta(t)$ is a CMV operator

$$(4.5) \quad \begin{aligned} L(t) &= \rho^-(t)\rho(t)\delta_{\text{even}}S^{--} + (\beta^-(t)\rho(t)\delta_{\text{even}} - \alpha^+(t)\rho(t)\delta_{\text{odd}})S^- - \beta(t)\alpha^+(t) \\ &+ (\beta(t)\rho^+(t)\delta_{\text{even}} - \alpha^{++}(t)\rho^+(t)\delta_{\text{odd}})S^+ + \rho^+(t)\rho^{++}(t)\delta_{\text{odd}}S^{++}, \end{aligned}$$

acting on sequences over \mathbb{Z} , where δ_{even} and δ_{odd} denote the characteristic functions of the even, odd integers,

$$(4.6) \quad \delta_{\text{even}} = \chi_{2\mathbb{Z}}, \quad \delta_{\text{odd}} = 1 - \delta_{\text{even}} = \chi_{2\mathbb{Z}+1},$$

respectively. Next, consider

$$(4.7) \quad \begin{aligned} P_{\underline{p}}(t) &= \frac{i}{2} \sum_{\ell=1}^{p_+} c_{p_+-\ell,+} ([L^\ell(t)]_+ - [L^\ell(t)]_-) \\ &- \frac{i}{2} \sum_{\ell=1}^{p_-} c_{p_--\ell,-} ([L^{-\ell}(t)]_+ - [L^{-\ell}(t)]_-) - \frac{i}{2} c_{\underline{p}} Q_d, \quad \underline{p} \in \mathbb{N}_0^2, \end{aligned}$$

with Q_d denoting the doubly infinite diagonal matrix

$$(4.8) \quad Q_d = ((-1)^k \delta_{k,\ell})_{k,\ell \in \mathbb{Z}}.$$

Then the AL hierarchy is equivalent to the Lax equation

$$(4.9) \quad \frac{d}{dt} L(t) - [P_{\underline{p}}(t), L(t)] = 0, \quad t \in \mathbb{R}.$$

To find an explicit expression we introduce

$$(4.10) \quad \begin{aligned} f_{\ell,\pm}(t) &= \sum_{k=0}^{\ell} c_{\ell-k,\pm} \hat{f}_{k,\pm}(t), & g_{\ell,\pm}(t) &= \sum_{k=0}^{\ell} c_{\ell-k,\pm} \hat{g}_{k,\pm}(t), \\ h_{\ell,\pm}(t) &= \sum_{k=0}^{\ell} c_{\ell-k,\pm} \hat{h}_{k,\pm}(t), \end{aligned}$$

where

$$(4.11) \quad \begin{aligned} \hat{f}_{\ell,+}(n,t) &= \alpha(n,t) \langle \delta_n, L^{\ell+1} \delta_n \rangle + \rho(n,t) \begin{cases} \langle \delta_{n-1}, L^{\ell+1}(t) \delta_n \rangle, & n \text{ even}, \\ \langle \delta_n, L^{\ell+1}(t) \delta_{n-1} \rangle, & n \text{ odd}, \end{cases} \\ \hat{f}_{\ell,-}(n,t) &= \alpha(n,t) \langle \delta_n, L^{-\ell} \delta_n \rangle + \rho(n,t) \begin{cases} \langle \delta_{n-1}, L^{-\ell}(t) \delta_n \rangle, & n \text{ even}, \\ \langle \delta_n, L^{-\ell}(t) \delta_{n-1} \rangle, & n \text{ odd}, \end{cases} \\ \hat{g}_{0,\pm} &= 1/2, & \hat{g}_{\ell,\pm}(n,t) &= \langle \delta_n, L^{\pm\ell}(t) \delta_n \rangle, \\ \hat{h}_{\ell,+}(n,t) &= \beta(n,t) \langle \delta_n, L^{\ell}(t) \delta_n \rangle + \rho(n,t) \begin{cases} \langle \delta_n, L^{\ell}(t) \delta_{n-1} \rangle, & n \text{ even}, \\ \langle \delta_{n-1}, L^{\ell}(t) \delta_n \rangle, & n \text{ odd}, \end{cases} \\ \hat{h}_{\ell,-}(n,t) &= \beta(n,t) \langle \delta_n, L^{-\ell-1}(t) \delta_n \rangle + \rho(n,t) \begin{cases} \langle \delta_n, L^{-\ell-1}(t) \delta_{n-1} \rangle, & n \text{ even}, \\ \langle \delta_{n-1}, L^{-\ell-1}(t) \delta_n \rangle, & n \text{ odd}. \end{cases} \end{aligned}$$

Then the \underline{p} th equation, $\underline{p} = (p_-, p_+) \in \mathbb{N}_0^2$, in the AL hierarchy is given by

$$(4.12) \quad \begin{aligned} \text{AL}_{\underline{p}}(\alpha, \beta) &= \begin{pmatrix} -i\dot{\alpha}(t) - \alpha(g_{p_+,+}(t) + g_{p_-,-}^-(t)) + f_{p_+-1,+}(t) - f_{p_- -1,-}^-(t) \\ -i\dot{\beta}(t) + \beta(g_{p_+,+}^-(t) + g_{p_-,-}(t)) - h_{p_+-1,-}(t) + h_{p_- -1,+}^-(t) \end{pmatrix} = 0, \\ \underline{p} &= (p_-, p_+) \in \mathbb{N}_0^2. \end{aligned}$$

Theorem 4.2. Fix some $\underline{p} = (p_+, p_-) \in \mathbb{N}_0^2$ such that $p_- = p_+ > 0$ and set $p = p_+ + p_- - 1$. Assume that $\alpha_0(n, t), \beta_0(n, t)$, with $\rho_0(n, t) \neq 0$, and $\alpha(n, t), \beta(n, t)$ are solutions of some equation in the Toda hierarchy $\text{AL}_{\underline{p}}$ such that there is one $n_0 \in \mathbb{Z}$ and two times $t_0 < t_1$ such that

$$(4.13) \quad \alpha_0(n_0 + j, t) = \alpha(n_0 + j, t), \quad \beta_0(n_0 + j, t) = \beta(n_0 + j, t), \quad 0 \leq j \leq p,$$

for $t \in (t_0, t_1)$. Then

$$(4.14) \quad \alpha_0(n, t) = \alpha(n, t), \quad \beta_0(n, t) = \beta(n, t)$$

for all $(n, t) \in \mathbb{Z} \times \mathbb{R}$.

Proof. Again drop the dependence on t for notational simplicity during this proof and we will use the same conventions as in the proof of Theorem 3.1.

The homogeneous quantities $\hat{f}_{\ell,\pm}$, $\hat{g}_{\ell,\pm}$, $\hat{h}_{\ell,\pm}$ are uniquely defined by the following recursion relations [9, Lem. C.5]:

$$\begin{aligned}\hat{g}_{0,+} &= \frac{1}{2}, \quad \hat{f}_{0,+} = -\alpha^+, \quad \hat{h}_{0,+} = \beta, \\ \hat{g}_{l+1,+} &= \sum_{k=0}^l \hat{f}_{l-k,+} \hat{h}_{k,+} - \sum_{k=1}^l \hat{g}_{l+1-k,+} \hat{g}_{k,+}, \\ \hat{f}_{l+1,+} &= \hat{f}_{l,+} - \alpha(\hat{g}_{l+1,+} + \hat{g}_{l+1,+}^-), \\ \hat{h}_{l+1,+} &= \hat{h}_{l,+} + \beta(\hat{g}_{l+1,+} + \hat{g}_{l+1,+}^-),\end{aligned}$$

and

$$\begin{aligned}\hat{g}_{0,-} &= \frac{1}{2}, \quad \hat{f}_{0,-} = \alpha, \quad \hat{h}_{0,-} = -\beta^+, \\ \hat{g}_{l+1,-} &= \sum_{k=0}^l \hat{f}_{l-k,-} \hat{h}_{k,-} - \sum_{k=1}^l \hat{g}_{l+1-k,-} \hat{g}_{k,-}, \\ \hat{f}_{l+1,-} &= \hat{f}_{l,-} + \alpha(\hat{g}_{l+1,-} + \hat{g}_{l+1,-}^-), \\ \hat{h}_{l+1,-} &= \hat{h}_{l,-} - \beta(\hat{g}_{l+1,-} + \hat{g}_{l+1,-}^-).\end{aligned}$$

From them we obtain

$$\begin{aligned}(4.15) \quad \hat{f}_{j,+}(n) &= - \left(\prod_{l=1}^j \rho(n+l)^2 \right) \alpha(n+j+1) + R(n+j, n-j+2) \\ &\quad + \left(\prod_{l=0}^{j-2} \rho(n-l)^2 \right) \alpha(n+1)^2 \beta(n-j+1), \\ (4.16) \quad \hat{f}_{j,-}(n) &= - \left(\prod_{l=1}^{j-1} \rho(n+l)^2 \right) \alpha(n)^2 \beta(n+j) + R(n+j-1, n-j+1) \\ &\quad + \left(\prod_{l=0}^{j-1} \rho(n-l)^2 \right) \alpha(n-j),\end{aligned}$$

$$\begin{aligned}(4.17) \quad \hat{g}_{j,+}(n) &= - \left(\prod_{l=1}^{j-1} \rho(n+l)^2 \right) \beta(n) \alpha(n+j) + R(n+j-1, n-j+2) \\ &\quad - \left(\prod_{l=0}^{j-2} \rho(n-l)^2 \right) \alpha(n+1) \beta(n-j+1)\end{aligned}$$

$$\begin{aligned}(4.18) \quad \hat{g}_{j,-}(n) &= - \left(\prod_{l=1}^{j-1} \rho(n+l)^2 \right) \alpha(n) \beta(n+j) + R(n+j-1, n-j+2) \\ &\quad - \left(\prod_{l=0}^{j-2} \rho(n-l)^2 \right) \beta(n+1) \alpha(n-j+1)\end{aligned}$$

$$\begin{aligned}
 \hat{h}_{j,+}(n) = & - \left(\prod_{l=1}^{j-1} \rho(n+l)^2 \right) \beta(n)^2 \alpha(n+j) + R(n+j-1, n-j+1) \\
 & + \left(\prod_{l=0}^{j-1} \rho(n-l)^2 \right) \beta(n-j),
 \end{aligned}
 \tag{4.19}$$

$$\begin{aligned}
 \hat{h}_{j,-}(n) = & - \left(\prod_{l=1}^j \rho(n+l)^2 \right) \beta(n+j+1) + R(n+j, n-j+2) \\
 & + \left(\prod_{l=0}^{j-2} \rho(n-l)^2 \right) \beta(n+1)^2 \alpha(n-j+1)
 \end{aligned}
 \tag{4.20}$$

for $j \in \mathbb{N}$. Note that it suffices to verify the $+$ case since the $-$ case follows from $\hat{f}_{j,\pm}(\alpha, \beta) = \hat{h}_{j,\pm}(\alpha, \beta)$ and $\hat{g}_{j,+}(\alpha, \beta) = \hat{g}_{j,-}(\alpha, \beta)$ ([9, Lem. 3.7]).

Now we can proceed as in the case of the Toda hierarchy. For example

$$0 = i(\dot{\alpha}(n+p_-) - \dot{\alpha}_0(n+p_-)) = -c_{0,+} \left(\prod_{l=p_-}^p \rho(n-l)^2 \right) (\alpha(n+p+1) - \alpha_0(n+p+1))$$

implies $\alpha(n+p+1) = \alpha_0(n+p+1)$ etc. \square

Interestingly, the above approach does not seem to work for $p_- \neq p_+$ in general. In any case, the above result covers the discrete nonlinear Schrödinger and Schur hierarchies via the above mentioned special choices $c_{0,\pm} = 1$, $\beta = \pm \bar{\alpha}$ and $c_{0,\pm} = \pm i$, $\beta = \bar{\alpha}$.

Acknowledgments. We thank F. Gesztesy for pointing out errors in a previous version of this article.

REFERENCES

- [1] W. Bulla, F. Gesztesy, H. Holden, and G. Teschl, *Algebro-Geometric Quasi-Periodic Finite-Gap Solutions of the Toda and Kac-van Moerbeke Hierarchies*, Mem. Amer. Math. Soc. **135-641** (1998).
- [2] J. Bourgain, *On the compactness of the support of solutions of dispersive equations*, Internat. Math. Res. Notices **1997**, no. 9, 437–447 (1997).
- [3] J. Bourgain, *Anderson-Bernoulli models*, Mosc. Math. J., **5:3**, 523–536 (2005).
- [4] J. Bourgain, C. E. Kenig, *On localization in the continuous Anderson-Bernoulli model in higher dimension*, Invent. Math. **161:2**, 389–426 (2005).
- [5] L. Escauriaza, C. E. Kenig, G. Ponce and L. Vega, *On uniqueness properties of solutions of Schrödinger equations*, Comm. Partial Differential Equations **31:12**, 1811–1823 (2006).
- [6] L. Faddeev and L. Takhtajan, *Hamiltonian Methods in the Theory of Solitons*, Springer, Berlin, 1987.
- [7] H. Flaschka, *The Toda lattice. I. Existence of integrals*, Phys. Rev. B **9**, 1924–1925 (1974).
- [8] F. Gesztesy, F. Gesztesy, H. Holden, J. Michor, and G. Teschl, *The Ablowitz–Ladik hierarchy revisited*, in “Methods of Spectral Analysis in Mathematical Physics”, J. Janas (ed.) et al., 139–190, Oper. Theory Adv. Appl. **186**, Birkhäuser, Basel, 2009.
- [9] F. Gesztesy, H. Holden, J. Michor, and G. Teschl, *Soliton Equations and Their Algebro-Geometric Solutions. Volume II: (1 + 1)-Dimensional Discrete Models*, Cambridge Studies in Advanced Mathematics **114**, Cambridge University Press, Cambridge, 2008.
- [10] A. Ionescu and C. E. Kenig, *L^p Carleman inequalities and uniqueness of solutions of nonlinear Schrödinger equations*, Acta Math. **193:2**, 193–239 (2004).
- [11] C. E. Kenig, G. Ponce and L. Vega, *On the support of solutions to the generalized KdV equation*, Ann. I. H. Poincaré **19:2**, 191–208 (2002).

- [12] C. E. Kenig, G. Ponce, L. Vega, *On unique continuation of solutions to the generalized KdV equation*, Math. Res. Lett. **10**, 833–846 (2003).
- [13] H. Krüger and G. Teschl, *Long-time asymptotics for the Toda lattice for decaying initial data revisited*, Rev. Math. Phys. **21:1**, 61–109 (2009).
- [14] J. Lenells, *Infinite propagation speed of the Camassa–Holm equation*, J. Math. Anal. Appl. **325:2**, 1468–1478 (2007).
- [15] J. Michor and G. Teschl, *On the equivalence of different Lax pairs for the Kac–van Moerbeke hierarchy*, in Modern Analysis and Applications, V. Adamyan (ed.) et al., 437–445, Oper. Theory Adv. Appl. **191**, Birkhäuser, Basel, 2009.
- [16] G. Teschl, *Jacobi Operators and Completely Integrable Nonlinear Lattices*, Math. Surv. and Mon. **72**, Amer. Math. Soc., Rhode Island, 2000.
- [17] G. Teschl, *Almost everything you always wanted to know about the Toda equation*, Jahresber. Deutsch. Math.-Verein. **103**, no. 4, 149–162 (2001).
- [18] G. Teschl, *On the spatial asymptotics of solutions of the Toda lattice*, arXiv:0901.2717.
- [19] M. Toda, *Theory of Nonlinear Lattices*, 2nd enl. ed., Springer, Berlin, 1989.
- [20] B. Zhang, *Unique continuation for the Korteweg–de Vries equation*, SIAM J. Math. Anal. **23**, 55–71 (1992).

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX 77005, USA

E-mail address: helge.krueger@rice.edu

URL: <http://math.rice.edu/~hk7/>

FACULTY OF MATHEMATICS, NORDBERGSTRASSE 15, 1090 WIEN, AUSTRIA, AND INTERNATIONAL ERWIN SCHRÖDINGER INSTITUTE FOR MATHEMATICAL PHYSICS, BOLTZMANNGASSE 9, 1090 WIEN, AUSTRIA

E-mail address: Gerald.Teschl@univie.ac.at

URL: <http://www.mat.univie.ac.at/~gerald/>