


# COMPLEX ANALYSIS

Lecture notes for  
MA 648

Rudi Weikard



Version of December 8, 2016



# Contents

Preface	1
Chapter 1. The complex numbers: algebra, geometry, and topology	3
1.1. The algebra of complex numbers	3
1.2. The topology and geometry of complex numbers	4
1.3. Sequences and series	5
Chapter 2. Complex-valued functions of a complex variable	7
2.1. Limits and continuity	7
2.2. Holomorphic functions	7
2.3. Integration	8
2.4. Series of functions	10
2.5. Analytic functions	11
Chapter 3. Cauchy's theorem and some of its consequences	13
3.1. The index of a point with respect to a closed contour	13
3.2. Cauchy's theorem	13
3.3. Consequences of Cauchy's theorem	14
3.4. The global version of Cauchy's theorem	15
Chapter 4. Isolated singularities	17
4.1. Classifying isolated singularities	17
4.2. The calculus of residues	18
Chapter 5. A zoo of functions	21
5.1. Polynomial functions	21
5.2. Rational functions	22
5.3. Exponential and trigonometric functions	22
5.4. The logarithmic function and powers	23
Chapter 6. Entire functions	27
6.1. Infinite products	27
6.2. Weierstrass's factorization theorem	27
6.3. Counting zeros using Jensen's theorem	28
6.4. Hadamard's factorization theorem	29
Appendix A. Topology in metric spaces	31
A.1. Basics	31
A.2. Compactness	32
A.3. Product spaces	32

Glossary	33
List of Symbols	35
Hints	37
Index	39

## Preface

Complex Analysis, also called the Theory of Functions, is one of the most important and certainly one of the most beautiful branches of mathematics. This is due to the fact that, in the case of complex variables, differentiability in open sets has consequences which are much more significant than in the case of real variables. It is the goal of this course to study these consequences and some of their far reaching applications.

As a prerequisite of the course familiarity with Advanced Calculus<sup>1</sup> is necessary and, generally, sufficient. Some prior exposure to topology may be useful but the necessary concepts and theorems are provided in the appendix if they are not covered in the course.

There are many textbooks on our subject; several of the following have been consulted in the preparation of these notes.

- Lars V. Ahlfors, *Complex analysis*, McGraw-Hill, New York, 1953.
- John B. Conway, *Functions of one complex variable*, Springer, New York, 1978.
- Walter Rudin, *Real and complex analysis*, McGraw-Hill, New York, 1987.
- Elias M. Stein and Rami Shakarchi, *Complex analysis*, Princeton University Press, Princeton, 2003.

In order to help you navigate the notes a glossary of terms, a list of symbols, and an index of central terms are appended at the end of these notes. Terms in the glossary are defined there and not in the text as they are not central, only convenient. The index points you to the places where terms central to Complex Analysis are defined in the notes.

These notes are not intended to be encyclopedic; I tried, rather, to be pedagogic. As a consequence there are many occasions where assumptions could be weakened or conclusions strengthened. Also, very many interesting results living close by have been skipped and other interesting and fruitful subjects have not even been touched. This is just the very beginning of the most exciting and beautiful Theory of Functions.

Thanks to my classes of Fall 2013 and Fall 2016 who caused many improvements to the notes and found a large number of mistakes (but probably not all).

---

<sup>1</sup>Among many sources my Advanced Calculus notes would serve in this respect. They may be found at <http://people.cas.uab.edu/~weikard/teaching/ac.pdf>.



## CHAPTER 1

# The complex numbers: algebra, geometry, and topology

### 1.1. The algebra of complex numbers

**1.1.1 The field of complex numbers.** The set  $\mathbb{C}$  of complex numbers is the set of all ordered pairs of real numbers equipped with two **binary operations**  $+$  and  $\cdot$  as follows: if  $(a, b)$  and  $(c, d)$  are pairs of real numbers, then

$$(a, b) + (c, d) = (a + c, b + d)$$

and

$$(a, b) \cdot (c, d) = (ac - bd, ad + bc).$$

The set  $\mathbb{C}$  with the binary operations  $+$  and  $\cdot$  is a field, i.e., the following properties hold:

- (A1):  $(x + y) + z = x + (y + z)$  for all  $x, y, z \in \mathbb{C}$  (*associative law of addition*).
- (A2):  $x + y = y + x$  for all  $x, y \in \mathbb{C}$  (*commutative law of addition*).
- (A3):  $(0, 0)$  is the unique additive **identity**.
- (A4): Each  $x \in \mathbb{C}$  has a unique additive **inverse**  $-x \in \mathbb{C}$ , called the *negative* of  $x$ .
- (M1):  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$  for all  $x, y, z \in \mathbb{C}$  (*associative law of multiplication*).
- (M2):  $x \cdot y = y \cdot x$  for all  $x, y \in \mathbb{C}$  (*commutative law of multiplication*).
- (M3):  $(1, 0)$  is the unique multiplicative **identity**.
- (M4): Each  $x \in \mathbb{C} \setminus \{(0, 0)\}$  has a unique multiplicative **inverse**  $x^{-1} \in \mathbb{C}$ , called the *reciprocal* of  $x$ .
- (D):  $(x + y) \cdot z = x \cdot z + y \cdot z$  for all  $x, y, z \in \mathbb{C}$  (*distributive law*).

For simplicity we will usually write  $x - y$  in place of  $x + (-y)$ , and  $\frac{x}{y}$  or  $x/y$  for  $x \cdot y^{-1}$ . It is also common to write  $xy$  instead of  $x \cdot y$  and to let multiplication take precedence over addition, i.e.,  $x + yz$  is short for  $x + (yz)$ .

If we identify the real number  $x$  with the complex number  $(x, 0)$  we find that addition and multiplication of real numbers are faithfully reproduced in  $\mathbb{C}$ : i.e.,  $(x, 0) + (y, 0) = (x + y, 0)$  and  $(x, 0)(y, 0) = (xy, 0)$ . Thus we may treat  $\mathbb{R}$  as a subfield of  $\mathbb{C}$ . Next note that  $(x, y) = (x, 0) + (0, 1)(y, 0)$ . Abbreviating  $(0, 1)$  by  $i$  this becomes  $x + iy$ , the standard notation for complex numbers. Numbers of the form  $iy$ ,  $y \in \mathbb{R}$ , are called (purely) *imaginary* numbers.

Complex numbers are not ordered. Still we will encounter inequalities like  $z \geq w$  and, if we do, we tacitly assume that  $z, w \in \mathbb{R}$ .

**1.1.2 Real and imaginary parts of a complex number.** If  $a, b \in \mathbb{R}$  and  $z = a + ib$  is a complex number, then  $a$  is called the *real part* of  $z$  (denoted by  $\text{Re}(z)$ ) and  $b$  is called the *imaginary part* of  $z$  (denoted by  $\text{Im}(z)$ ). The number  $\bar{z} = a - ib$  is called the *complex conjugate* of  $z$ . If  $z \in \mathbb{C}$ , then  $z + \bar{z} = 2\text{Re}(z)$ ,  $z - \bar{z} = 2i\text{Im}(z)$ , and  $z\bar{z} = \text{Re}(z)^2 + \text{Im}(z)^2 \geq 0$ . If  $z, w \in \mathbb{C}$ , then  $\overline{z + w} = \bar{z} + \bar{w}$  and  $\overline{z\bar{w}} = \bar{z}\bar{w}$ .

**1.1.3 Absolute value of a complex number.** If  $a, b \in \mathbb{R}$  and  $z = a + ib$  is a complex number, then  $|z| = \sqrt{a^2 + b^2} \in [0, \infty)$  is called the *absolute value* or *modulus* of  $z$ . Note that this definition is compatible with the definition of the absolute value of a real number.

If  $z \in \mathbb{C}$ , then  $|z|^2 = z\bar{z}$ ,  $|\operatorname{Re}(z)| \leq |z|$ , and  $|\operatorname{Im}(z)| \leq |z|$ . If  $z \neq 0$ , then  $1/z = \bar{z}/|z|^2$ . The absolute value of a product of two complex numbers equals the product of the absolute values of the numbers, i.e.,  $|z_1 z_2| = |z_1| |z_2|$ .

**1.1.4 The triangle inequality.** If  $x, y \in \mathbb{C}$ , then

$$|x + y| \leq |x| + |y| \quad \text{and} \quad |x + y| \geq ||x| - |y||.$$

## 1.2. The topology and geometry of complex numbers

**1.2.1 Distance between complex numbers.** Given two complex number  $z_1$  and  $z_2$  we call  $|z_1 - z_2|$  their *distance*. In fact, equipped with this distance function,  $\mathbb{C}$  becomes a metric space and thus an object of geometry. When one wants to emphasize the geometry of  $\mathbb{C}$  one calls it the *complex plane*. Appendix A gathers some pertinent information about metric (and topological) spaces. In particular, it reviews the concepts of convergence and limit in metric spaces, which are obvious generalizations of those in  $\mathbb{R}$ .

**1.2.2 Polar representation of complex numbers.** For any non-zero complex number  $z = x + iy$ ,  $x, y \in \mathbb{R}$  define  $r = \sqrt{x^2 + y^2} = |z| > 0$  and  $\theta \in (-\pi, \pi]$  by the relations  $\cos \theta = x/r$  and  $\sin \theta = y/r$ . These requirements determine  $r$  and  $\theta$  uniquely. Interpreting  $z$  as a point  $(x, y)$  in a two-dimensional rectangular coordinate system the number  $r$  represents the distance of  $z$  from the origin. If  $\theta \geq 0$ , it is the counterclockwise angle from the positive  $x$ -axis to the ray connecting  $z$  and the origin, otherwise  $-\theta$  is the clockwise angle. The numbers  $r$  and  $\theta$  are called the *polar coordinates* of  $z$ .

**1.2.3 Powers and roots.** For any  $z \in \mathbb{C}$  we define  $z^0 = 1$  and, inductively,  $z^n = z z^{n-1}$  for  $n \in \mathbb{N}$ . When  $z \neq 0$  and  $n \in \mathbb{N}$  we note that  $(z^{-1})^n = (z^n)^{-1}$  and we abbreviate this number by  $z^{-n}$ . The numbers  $z^n$ ,  $n \in \mathbb{Z}$ , are called the (integer) *powers* of  $z$ .

We also define *n-th roots* of complex numbers when  $n \in \mathbb{N}$ . The complex number  $a$  is called an *n-th root* of a complex number  $b$ , if  $a^n = b$ . Every non-zero complex number has exactly  $n$  *n-th roots*. [Hint](#)

**1.2.4 Open and closed sets.** As a metric space  $\mathbb{C}$  has open and closed balls which, in this context, are usually called disks. Thus, as explained in [A.1.4](#),  $\mathbb{C}$  is a topological space whose open sets are unions of open disks; in particular open disks are open. It is also true that closed disks are closed. The following characterizations of open and closed sets are often very useful.

- A set  $S \subset \mathbb{C}$  is open if and only if for every  $x \in S$  there is an  $r > 0$  such that  $B(x, r) \subset S$ .
- A set  $S \subset \mathbb{C}$  is closed if and only if for every convergent sequence whose elements lie in  $S$  the limit also lies in  $S$ .

As a metric space  $\mathbb{C}$  is the same as  $\mathbb{R}^2$ . Hence, by the Heine-Borel theorem [A.2.5](#), a subset of the complex plane is compact precisely when it is closed and bounded.

**1.2.5 Line segments and convex sets.** Given two points  $z$  and  $w$  in  $\mathbb{C}$  we call the set  $\{(1-t)z + tw : 0 \leq t \leq 1\}$  the *line segment* joining  $z$  and  $w$ . The number  $|z - w|$  is called the *length* of the segment.

A subset  $S$  of  $\mathbb{C}$  is called *convex* if it contains the line segment joining  $x$  and  $y$  whenever  $x, y \in S$ . Every disk is convex.



**1.2.6 Connectedness.** A subset  $S$  of  $\mathbb{C}$  is called *not connected* if there are two open sets  $A$  and  $B$  in  $\mathbb{C}$  such that  $S \subset A \cup B$  and  $S \cap A$  and  $S \cap B$  are non-empty disjoint sets. If no such sets exist, then  $S$  is called *connected*.

The empty set, all **singletons**, and all convex sets are connected. **Hint**

**1.2.7 Connected components.** If  $S$  is a subset of the complex plane and  $x \in S$ , let  $\mathcal{C}(x)$  be the collection of all connected subsets of  $S$  which contain  $x$ . Any union of sets in  $\mathcal{C}(x)$  is connected. In particular,  $T = \bigcup_{C \in \mathcal{C}(x)} C$  is a connected subset of  $S$ . Any larger subset of  $S$  is not connected.  $T$  is called a component of  $S$ . The components of  $S$  are pairwise disjoint and their union equals  $S$ .

The components of open sets are open. There are at most countably many of them.

### 1.3. Sequences and series

**1.3.1 Numerical sequences.** A sequence  $z$  of complex numbers converges if and only if both of the real sequences  $n \mapsto \operatorname{Re}(z_n)$  and  $n \mapsto \operatorname{Im}(z_n)$  converge. In this case

$$\lim_{n \rightarrow \infty} z_n = \lim_{n \rightarrow \infty} \operatorname{Re}(z_n) + i \lim_{n \rightarrow \infty} \operatorname{Im}(z_n).$$

This result allows to extend immediately many theorems proved for sequences of real numbers to sequences of complex numbers. In particular, if the sequences  $z$  and  $w$  have limits, then so do  $z \pm w$  and  $zw$ . In fact,

$$\lim_{n \rightarrow \infty} (z_n \pm w_n) = \lim_{n \rightarrow \infty} z_n \pm \lim_{n \rightarrow \infty} w_n,$$

$$\lim_{n \rightarrow \infty} z_n w_n = \lim_{n \rightarrow \infty} z_n \lim_{n \rightarrow \infty} w_n,$$

and, assuming also  $w_n \neq 0$  for all  $n$  and  $\lim_{n \rightarrow \infty} w_n \neq 0$ ,

$$\lim_{n \rightarrow \infty} \frac{z_n}{w_n} = \frac{\lim_{n \rightarrow \infty} z_n}{\lim_{n \rightarrow \infty} w_n}.$$

**1.3.2 Cauchy sequences.** A sequence  $z$  of complex numbers is called a *Cauchy*<sup>1</sup> sequence if

$$\forall \varepsilon > 0 : \exists N > 0 : \forall n, m > N : |z_n - z_m| < \varepsilon.$$

**THEOREM.** A sequence of complex numbers converges if and only if it is a Cauchy sequence.

**1.3.3 Series.** If  $z : \mathbb{N} \rightarrow \mathbb{C}$  is a sequence of complex numbers the sequence

$$s : \mathbb{N} \rightarrow \mathbb{C} : n \mapsto s_n = \sum_{k=1}^n z_k$$

is called the *sequence of partial sums* of  $z$  or a *series*. We will denote  $s$  by  $\sum_{k=1}^{\infty} z_k$ .

If the sequence  $s$  of partial sums of  $z$  converges to  $L \in \mathbb{C}$ , we say the series  $\sum_{k=1}^{\infty} z_k$  *converges* to  $L$ . We then write (abusing notation slightly)

$$\sum_{k=1}^{\infty} x_k = \lim_{n \rightarrow \infty} s_n = L.$$

Of course, a series (i.e., a sequence of partial sums) may also *diverge*.

**1.3.4 Absolute convergence of series.** The series  $\sum_{k=1}^{\infty} z_k$  is called *absolutely convergent* if  $\sum_{k=1}^{\infty} |z_k|$  is convergent. Every absolutely convergent series is convergent.

<sup>1</sup>Augustin-Louis Cauchy (1789 – 1857)

THEOREM. Suppose that  $\sum_{n=1}^{\infty} z_n$  converges absolutely and let  $\sum_{n=1}^{\infty} w_n$  be a **re-arrangement** of  $\sum_{n=1}^{\infty} z_n$ . Then  $\sum_{n=1}^{\infty} w_n$  converges absolutely, and

$$\sum_{n=1}^{\infty} w_n = \sum_{n=1}^{\infty} z_n.$$

**Hint**

**1.3.5 The geometric series.** If  $|z| < 1$ , then the *geometric series*  $\sum_{n=0}^{\infty} z^n$  converges absolutely to  $1/(1 - z)$ .

## CHAPTER 2

# Complex-valued functions of a complex variable

### 2.1. Limits and continuity

**2.1.1 Limits.** Let  $S$  be a subset of  $\mathbb{C}$ ,  $f : S \rightarrow \mathbb{C}$  a function, and  $a \in \mathbb{C}$  a **limit point** of  $S$ . We say that  $f$  *converges* to the complex number  $b$  as  $z$  tends to  $a$  if the following statement is true:

$$\forall \varepsilon > 0 : \exists \delta > 0 : \forall z \in S : 0 < |z - a| < \delta \Rightarrow |f(z) - b| < \varepsilon.$$

Note that this definition can easily be extended to any metric space.

As for sequences,  $f$  can converge to at most one value as  $z$  tends to  $a$ . This value is then called the limit of  $f$  as  $z$  tends to  $a$ . It is denoted by  $\lim_{z \rightarrow a} f(z)$ .

The results on limits of sequences stated in **1.3.1** hold analogously also for functions. Moreover, if  $g$  has limit  $b$  at  $a$  and  $f$  has limit  $c$  at  $b$ , then  $f \circ g$  has limit  $c$  at  $a$  (assuming, of course, that  $b$  and the range of  $g$  are in the domain of  $f$ ).

**2.1.2 Continuity.** Let  $S$  be a subset of  $\mathbb{C}$  and  $a \in S$ . A function  $f : S \rightarrow \mathbb{C}$  is continuous at  $a$  if the following statement is true:

$$\forall \varepsilon > 0 : \exists \delta > 0 : \forall z \in S : |z - a| < \delta \Rightarrow |f(z) - f(a)| < \varepsilon.$$

Any function is continuous at any **isolated point** of its domain. Suppose  $S \subset \mathbb{C}$ ,  $f : S \rightarrow \mathbb{C}$ , and  $a \in S$  is a limit point of  $S$ . Then  $f$  is continuous at  $a$  if and only if  $\lim_{z \rightarrow a} f(z)$  exists and equals  $f(a)$ . It follows that sums and products and compositions of continuous functions are continuous.

### 2.2. Holomorphic functions

**2.2.1 Differentiation.** Let  $S$  be a subset of  $\mathbb{C}$ ,  $f : S \rightarrow \mathbb{C}$  a function, and  $a \in S$  a limit point of  $S$ . We say that  $f$  is *differentiable* at  $a$  if

$$\lim_{z \rightarrow a} \frac{f(z) - f(a)}{z - a}$$

exists. The limit is called the *derivative* of  $f$  at  $a$  and is commonly denoted by  $f'(a)$ . If  $f : S \rightarrow \mathbb{C}$  is differentiable at every point  $a \in S$ , we say that  $f$  is differentiable on  $S$ .

This definition is completely analogous to that of a derivative of a function of a real variable. In fact, when  $S$  is a real interval (one of the cases we are interested in) things are hardly different from Real Analysis; the fact that  $f$  assumes complex values is not very important. However, if  $S$  is an open set and  $z$  may approach  $a$  from many directions, the existence of the limit has far-reaching consequences.

**2.2.2 Basic properties of derivatives.** Suppose  $S$ ,  $S_1$  and  $S_2$  are subsets of  $\mathbb{C}$ . Then the following statements hold true:

- (1) A function  $f : S \rightarrow \mathbb{C}$  is differentiable at a point  $a \in S$  if and only if there is a number  $F$  and a function  $h : S \rightarrow \mathbb{C}$  which is continuous at  $a$ , vanishes there, and satisfies  $f(z) = f(a) + F(z - a) + h(z)(z - a)$ .
- (2) If  $f : S \rightarrow \mathbb{C}$  is differentiable at  $a \in S$ , then it is continuous at  $a$ .
- (3) Sums and products of differentiable functions with common domains are again differentiable.
- (4) The chain rule holds, i.e., if  $f : S_1 \rightarrow S_2$  is differentiable at  $a$  and  $g : S_2 \rightarrow \mathbb{C}$  is differentiable at  $f(a)$ , then  $g \circ f$  is differentiable at  $a$ , and  $(g \circ f)'(a) = g'(f(a))f'(a)$ .

Hint

**2.2.3 Differentiation with respect to real and complex variables.** Suppose  $S$  is a real interval. Then the function  $f : S \rightarrow \mathbb{C}$  is differentiable at  $a \in S$  if and only if  $\operatorname{Re}(f)$  and  $\operatorname{Im}(f)$  are differentiable there. In this case,  $f' = \operatorname{Re}(f)' + i \operatorname{Im}(f)'$ .

The situation is completely different if  $S$  is an open set. For instance, the function  $z \mapsto f(z) = z^2$  is differentiable on all of  $\mathbb{C}$  and  $f'(z) = 2z$ . However, its real and imaginary parts are differentiable only at  $z = 0$ . In fact, if a real-valued function defined in  $B(a, r)$  is differentiable at  $a$ , then its derivative must be zero.

**2.2.4 Holomorphic functions.** Let  $\Omega$  be a non-empty open set. A function  $f : \Omega \rightarrow \mathbb{C}$  is called *holomorphic* on  $\Omega$  if it is differentiable at every point of  $\Omega$ . A function which is defined and holomorphic on all of  $\mathbb{C}$  is called *entire*.

It is easy to think that the notions of differentiability and holomorphicity are the same. To avoid this mistake note that differentiability is a pointwise concept while holomorphicity is defined on open sets where every point is at the center of a disk contained in the set.

**2.2.5 Basic properties of holomorphic functions.** Holomorphic functions are continuous on their domain.

Sums, differences, and products of holomorphic functions (on common domains) are holomorphic. The composition of holomorphic functions is also holomorphic. The usual formulas hold, including the chain rule.

All polynomials (of a single variable) are entire functions. The function  $z \mapsto 1/z$  is holomorphic on  $\mathbb{C} \setminus \{0\}$ . The quotient  $f/g$  of two holomorphic functions  $f, g : \Omega \rightarrow \mathbb{C}$  is holomorphic on  $\Omega \setminus \{z \in \Omega : g(z) = 0\}$ .

## 2.3. Integration

**2.3.1 Paths.** Let  $\Omega \subset \mathbb{C}$  be a non-empty open set and  $[a, b]$  a non-trivial bounded interval in  $\mathbb{R}$ . A continuous function  $\gamma : [a, b] \rightarrow \Omega$  is then called a *path* in  $\Omega$ . If the derivative  $\gamma'$  exists and is continuous on  $[a, b]$ ,  $\gamma$  is called a *smooth path*.

The points  $\gamma(a)$  and  $\gamma(b)$  are called *initial point* and *end point* of  $\gamma$ , respectively. A path is called closed if its initial and end points coincide. The range of  $\gamma$ , i.e., the set  $\{\gamma(t) : t \in [a, b]\}$  is denoted by  $\gamma^*$ . Note that  $\gamma^*$  is compact and connected.

The path  $\eta : [a, b] \rightarrow \Omega$  is called the *opposite* of  $\gamma : [a, b] \rightarrow \Omega$  if  $\eta(t) = \gamma(a + b - t)$ . We denote the opposite of  $\gamma$  by  $\ominus\gamma$ .

The number  $\int_a^b |\gamma'(t)| dt$  is called the *length* of the smooth path  $\gamma$ .

**2.3.2 Reparametrization.** The smooth path  $\gamma_2 : [a_2, b_2] \rightarrow \Omega$  is called equivalent to the smooth path  $\gamma_1 : [a_1, b_1] \rightarrow \Omega$  if there exists a strictly increasing, continuously differentiable, and surjective function  $\varphi : [a_1, b_1] \rightarrow [a_2, b_2]$  such that  $\gamma_2 \circ \varphi = \gamma_1$ . Equivalence of smooth paths is an equivalence relation, i.e., it has the following three properties:

- (1) Any smooth path is equivalent to itself (reflexivity).

- (2) If  $\gamma_2$  is equivalent to  $\gamma_1$ , then  $\gamma_1$  is equivalent to  $\gamma_2$  (symmetry).  
 (3) If  $\gamma_2$  is equivalent to  $\gamma_1$  and  $\gamma_3$  is equivalent to  $\gamma_2$ , then  $\gamma_3$  is equivalent to  $\gamma_1$  (transitivity).

Given a smooth path it is always possible to find an equivalent one whose domain is  $[0, 1]$ .

**2.3.3 Integrals of complex-valued functions over intervals.** Let  $[a, b]$  be a closed interval in  $\mathbb{R}$  and  $f$  a complex-valued function on  $[a, b]$ . Then  $\operatorname{Re}(f)$  and  $\operatorname{Im}(f)$  are real-valued functions on  $[a, b]$ . We say that  $f$  is Riemann integrable over  $[a, b]$  if  $\operatorname{Re}(f)$  and  $\operatorname{Im}(f)$  are. The integral is defined to be

$$\int_a^b f = \int_a^b \operatorname{Re}(f) + i \int_a^b \operatorname{Im}(f).$$

The integral is **linear** and satisfies

$$\left| \int_a^b f \right| \leq \int_a^b |f|.$$

**Hint**

**2.3.4 Integrals along smooth paths.** Suppose  $\gamma : [a, b] \rightarrow \mathbb{C}$  is a smooth path in  $\mathbb{C}$  and  $f : \gamma^* \rightarrow \mathbb{C}$  is continuous. Then the number

$$\int_a^b (f \circ \gamma) \gamma'$$

is well-defined. It is called the integral of  $f$  along  $\gamma$  and denoted by  $\int_\gamma f$ . We have the estimate

$$\left| \int_\gamma f \right| \leq \sup(|f|(\gamma^*)) L(\gamma)$$

where  $L(\gamma)$  is the length of  $\gamma$ .

The integrals of  $f$  along equivalent smooth paths are all identical. The integral  $\int_{-\gamma} f$  along the opposite of a smooth path  $\gamma$  is the negative of  $\int_\gamma f$ .

**2.3.5 Example.** Let  $\gamma : [0, 2\pi] \rightarrow \mathbb{C} : t \mapsto \cos(t) + i \sin(t)$  and  $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C} : z \mapsto 1/z$ . Then  $\int_\gamma f = 2\pi i$ .

**2.3.6 Contours.** Let  $\Omega$  be a non-empty open set. Given several (not necessarily distinct) smooth paths in  $\Omega$ , say  $\gamma_1, \dots, \gamma_n$ , we may be interested in integrating along them successively. We call such a collection of smooth paths in  $\Omega$  (a **multiset**, actually, since we allow repetition) a *contour* in  $\Omega$  and write  $\Gamma = \gamma_1 \oplus \dots \oplus \gamma_n$ . Then we define

$$\int_\Gamma f = \sum_{k=1}^n \int_{\gamma_k} f.$$

The set  $\bigcup_{k=1}^n \gamma_k^*$  is denoted by  $\Gamma^*$ . It is a compact set.

Of course, we will write  $\gamma_1 \ominus \gamma_2$  for  $\gamma_1 \oplus (\ominus \gamma_2)$ .

Suppose  $\gamma : [a, b] \rightarrow \mathbb{C}$  is a smooth path in  $\mathbb{C}$  and  $f : \gamma^* \rightarrow \mathbb{C}$  is continuous. If  $c \in (a, b)$  and  $\gamma_1$  and  $\gamma_2$  are the restrictions of  $\gamma$  to  $[a, c]$  and  $[c, b]$ , respectively, then  $\int_\gamma f = \int_{\gamma_1 \oplus \gamma_2} f$ .

**2.3.7 Closed contours.** A contour  $\gamma_1 \oplus \dots \oplus \gamma_n$  is called *closed* when there is a **permutation**  $\pi$  of  $\{1, \dots, n\}$  such that the end point of  $\gamma_k$  coincides with the initial point of  $\gamma_{\pi(k)}$  for  $k = 1, \dots, n$ .

**2.3.8 Connected contours.** A particularly important instance of a contour  $\Gamma = \gamma_1 \oplus \dots \oplus \gamma_n$  is when for  $k = 1, \dots, n-1$ , the end point of  $\gamma_k$  coincides with the initial point of  $\gamma_{k+1}$  (perhaps after reordering the indices). A contour of this type is called a *connected contour*. The initial point of  $\gamma_1$  is called the *initial point* of  $\Gamma$  while the end point of  $\gamma_n$  is called the *end point* of  $\Gamma$ . We say a contour connects  $x$  to  $y$  if it is a connected contour with initial point  $x$  and end point  $y$ .

If  $\Gamma$  is a connected contour, then  $\Gamma^*$  is connected.

**THEOREM.** Let  $S$  be a subset of  $\mathbb{C}$ . If any two points of  $S$  are connected by a connected contour in  $S$ , then  $S$  is connected. Conversely, if  $S$  is open and connected and  $x, y$  are two points in  $S$ , then there is a connected contour in  $S$  which connects  $x$  and  $y$ .

**Hint**

**2.3.9 Polygonal contours.** Let  $z_1, \dots, z_{n+1} \in \mathbb{C}$ . We define  $\gamma_k(t) = (1-t)z_k + tz_{k+1}$  for  $t \in [0, 1]$  and  $k = 1, \dots, n$ . Then  $\gamma_k$  is a smooth path with initial point  $z_k$  and end point  $z_{k+1}$ . Note that  $\gamma_k^*$  is the line segment joining  $z_k$  and  $z_{k+1}$ . The length of the line segment equals the length of  $\gamma_k$ . Moreover,  $\gamma_1 \oplus \dots \oplus \gamma_n$  is a connected contour which we denote by  $\langle z_1, \dots, z_{n+1} \rangle$ . In particular, when  $n = 3$  and  $z_4 = z_1$ ,  $\Gamma = \langle z_1, z_2, z_3, z_1 \rangle$  is a closed contour tracing the circumference of a (possibly degenerate) triangle.

Suppose  $z'_1 \in \langle z_2, z_3 \rangle^*$ ,  $z'_2 \in \langle z_3, z_1 \rangle^*$ , and  $z'_3 \in \langle z_1, z_2 \rangle^*$  and  $\Lambda = \langle z_1, z'_3, z'_2, z_1 \rangle \oplus \langle z_2, z'_1, z'_3, z_2 \rangle \oplus \langle z_3, z'_2, z'_1, z_3 \rangle \oplus \langle z'_3, z'_1, z'_2, z'_3 \rangle$ . If  $f$  is continuous on  $\Gamma^*$ , then  $\int_\Gamma f = \int_\Lambda f$ .

**2.3.10 Primitives.** Suppose  $\Omega$  is a non-empty open set and  $F' = f$  on  $\Omega$ . Then  $F$  is called a *primitive* of  $f$ . If  $F$  is a primitive of  $f$  and  $c$  is a constant, then  $F + c$  is also a primitive of  $f$ .

If  $\gamma : [a, b] \rightarrow \Omega$  is a smooth path and the continuous function  $f$  has a primitive  $F$  in  $\Omega$ , then  $\int_\gamma f = F(\gamma(b)) - F(\gamma(a))$ .

The primitives of the zero function in a non-empty connected open set are precisely the constant functions.

## 2.4. Series of functions

**2.4.1 Pointwise and uniform convergence.** Let  $S \subset \mathbb{C}$  and, for each  $n \in \mathbb{N}$ , let  $f_n$  be a function from  $S$  to  $\mathbb{C}$ . The map  $n \mapsto f_n$  is called a *sequence* of functions. We say that  $n \mapsto f_n$  *converges pointwise* to a function  $f : S \rightarrow \mathbb{C}$  if for *each* point  $z \in S$  the numerical sequence  $n \mapsto f_n(z)$  converges to  $f(z)$ . We say that  $n \mapsto f_n$  *converges uniformly* to a function  $f : S \rightarrow \mathbb{C}$  if for every  $\varepsilon > 0$  there exists  $N \in \mathbb{R}$  such that for all  $n > N$  and *all*  $z \in S$  we have that  $|f_n(z) - f(z)| < \varepsilon$ .

To reiterate, consider the statements

- (1)  $\forall z \in S : \forall \varepsilon > 0 : \exists N \in \mathbb{R} : \forall n > N : |f_n(z) - f(z)| < \varepsilon$ .
- (2)  $\forall \varepsilon > 0 : \exists N \in \mathbb{R} : \forall z \in S : \forall n > N : |f_n(z) - f(z)| < \varepsilon$ .

A sequence  $n \mapsto f_n$  converges to  $f$  pointwise if (1) holds and uniformly if (2) holds.

Obviously, uniform convergence implies pointwise convergence but not vice versa.

A series of functions is defined as the sequence of partial sums of functions with a common domain. Thus the definitions of pointwise and uniform convergence extend also to series of functions.

**2.4.2 Cauchy criterion for uniform convergence.** Let  $S \subset \mathbb{C}$  and  $f_n : S \rightarrow \mathbb{C}$  for  $n \in \mathbb{N}$ . The sequence of functions  $n \mapsto f_n$  converges uniformly on  $S$  if and only if for

all  $\varepsilon > 0$  there exists  $N \in \mathbb{R}$  such that for all  $n, m > N$  and all  $z \in S$  it holds that  $|f_n(z) - f_m(z)| < \varepsilon$ .

**2.4.3 The Weierstrass<sup>1</sup>  $M$ -test.** Let  $S \subset \mathbb{C}$  and suppose  $n \mapsto g_n$  is a sequence of complex-valued functions defined on  $S$ . Assume that there are non-negative numbers  $M_n$  such that  $|g_n(z)| \leq M_n$  for all  $z \in S$  and that the series  $\sum_{n=1}^{\infty} M_n$  converges. Then the following statements hold:

- (1) The series  $\sum_{n=1}^{\infty} g_n(z)$  converges absolutely for every  $z \in S$ .
- (2) The series  $\sum_{n=1}^{\infty} g_n$  converges uniformly in  $S$ .

**2.4.4 Uniform convergence and continuity.** Let  $S \subset \mathbb{C}$  and  $n \mapsto f_n$  a sequence of continuous complex-valued functions on  $S$  which converges uniformly to a function  $f : S \rightarrow \mathbb{C}$ . Then  $f$  is continuous on  $S$ .

**2.4.5 Uniform convergence and integration.** Let  $[a, b]$  be a bounded interval in  $\mathbb{R}$ . Suppose that  $n \mapsto f_n : [a, b] \rightarrow \mathbb{C}$  is a sequence of functions which are Riemann integrable over  $[a, b]$  and that this sequence converges uniformly to a function  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f$  is Riemann integrable and

$$\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n.$$

Of course, the analogous result holds for series: Suppose that  $n \mapsto g_n : [a, b] \rightarrow \mathbb{C}$  is a sequence of functions which are Riemann integrable over  $[a, b]$  and that the series  $\sum_{n=0}^{\infty} g_n$  converges uniformly to a function  $g : [a, b] \rightarrow \mathbb{C}$ . Then  $g$  is Riemann integrable and

$$\int_a^b g = \sum_{n=0}^{\infty} \int_a^b g_n.$$

## 2.5. Analytic functions

**2.5.1 Power Series.** Let  $z \in \mathbb{C}$  and  $a_n \in \mathbb{C}$  for all non-negative integers  $n$ . Then the series  $\sum_{n=0}^{\infty} a_n z^n$  is called a *power series*. If the series converges for every  $z \in \mathbb{C}$  we say that it has infinite radius of convergence. Otherwise there is a non-negative number  $R$  such that the series converges absolutely whenever  $|z| < R$  and diverges whenever  $|z| > R$ . The number  $R$  is called the *radius of convergence* of the series. The disk  $B(0, R)$  is called the disk of convergence.

The series  $\sum_{n=0}^{\infty} a_n z^n$  and  $\sum_{n=1}^{\infty} n a_n z^{n-1}$  have the same radius of convergence.

**2.5.2 Power series define holomorphic functions.** A power series with a positive or infinite radius of convergence defines a holomorphic function in its disk of convergence. In fact, if  $f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$  for  $|z - z_0| < R$ , then  $f'(z) = \sum_{n=1}^{\infty} n a_n (z - z_0)^{n-1}$  in the same disk, i.e., the power series may be differentiated term by term and  $f$  is holomorphic on  $B(z_0, R)$ .

**Hint**

**2.5.3 Analytic functions.** We say that a function  $f : B(z_0, R) \rightarrow \mathbb{C}$  is represented by a power series if  $f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$  for all  $z \in B(z_0, R)$ .

Let  $\Omega$  be a non-empty open set. A function  $f : \Omega \rightarrow \mathbb{C}$  is called *analytic* in  $\Omega$  if  $\Omega$  is a union of open disks in each of which  $f$  is represented by a power series.

For example,  $z \mapsto 1/(1 - z)$  is analytic in the (open) **unit disk** as well as in all of  $\mathbb{C} \setminus \{1\}$ .

---

<sup>1</sup>Karl Weierstraß (1815 – 1897)

THEOREM. Every analytic function is holomorphic.

The central and most astonishing theorem of complex analysis is that the converse is also true, i.e., every holomorphic function is analytic, see Theorem 3.3.2 below.

**2.5.4 Taylor series.** Suppose  $f : B(z_0, R) \rightarrow \mathbb{C}$  is analytic, i.e.,  $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ . Then  $f$  is infinitely often differentiable and  $a_n = f^{(n)}(z_0)/n!$ . We call

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n$$

the Taylor<sup>2</sup> series of  $f$  about  $z_0$ .

**2.5.5 Analytic functions defined by integrals.** Suppose  $\psi$  and  $\phi$  are continuous complex-valued functions on  $[a, b]$ . Then the function  $f$  defined by

$$f(z) = \int_a^b \frac{\psi(t)}{\phi(t) - z} dt$$

is analytic in  $\Omega = \mathbb{C} \setminus \phi([a, b])$ . In fact, if  $z_0 \in \Omega$ , if  $r = \inf\{|\phi(t) - z_0| : t \in [a, b]\}$ , and if  $z \in B(z_0, r)$ , then

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

where

$$a_n = \int_a^b \frac{\psi(t) dt}{(\phi(t) - z_0)^{n+1}}.$$

Hint

**2.5.6 The exponential function.** The series

$$\exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges absolutely for every  $z \in \mathbb{C}$  and uniformly in every closed disk about the origin. It represents therefore an entire function, called the *exponential function*. It is an extension of the exponential function defined in Real Analysis.

THEOREM. If  $t \in \mathbb{R}$ , then  $\exp(it) = \cos(t) + i \sin(t)$ ; in particular,  $\exp(0) = 1$ . Moreover  $\exp'(z) = \exp(z)$ , and  $\exp(a + b) = \exp(a) \exp(b)$ . The exponential function has no **zeros**. It is **periodic** with period  $2\pi i$ . Any **period** of the exponential function is an integer multiple of  $2\pi i$ . In particular,  $\exp(z) = 1$  if and only if  $z$  is an integer multiple of  $2\pi i$ .

---

<sup>2</sup>Brook Taylor (1685 – 1731)



## CHAPTER 3

# Cauchy's theorem and some of its consequences

### 3.1. The index of a point with respect to a closed contour

**3.1.1.** Let  $\Gamma$  be a contour in  $\mathbb{C}$  and  $\Omega = \mathbb{C} \setminus \Gamma^*$ . Since  $\Gamma^*$  is compact  $\Omega$  is open and contains the complement  $E$  of  $B(0, R)$  for some  $R > 0$ . The set  $E$  is connected and is part of precisely one connected component of  $\Omega$ .

**3.1.2.** If  $\gamma : [a, b] \rightarrow \mathbb{C}$  is a smooth path and  $z \in \mathbb{C} \setminus \gamma^*$ , define  $F : [a, b] \rightarrow \mathbb{C}$  by

$$F(s) = \exp \left( \int_a^s \frac{\gamma'(t)}{\gamma(t) - z} dt \right).$$

Then  $s \mapsto F(s)/(\gamma(s) - z)$  is constant.

**3.1.3 The index of a point with respect to a closed contour.** Let  $\Gamma$  be a closed contour in  $\mathbb{C}$  and  $\Omega = \mathbb{C} \setminus \Gamma^*$ . The function  $\text{Ind}_\Gamma$  defined by

$$\text{Ind}_\Gamma(z) = \frac{1}{2\pi i} \int_\Gamma \frac{du}{u - z}$$

on  $\Omega$  is an analytic function assuming only integer values. It is constant on each connected component of  $\Omega$  and, in particular, zero near infinity.

**3.1.4 Example.** Consider the smooth paths  $\gamma_1(t) = \exp(it)$ ,  $\gamma_2(t) = \exp(it)/2$ , and  $\gamma_3(t) = \exp(-it)/2$  all defined on  $[0, 2\pi]$  and the contours  $\Gamma_1 = \gamma_1$ ,  $\Gamma_2 = \gamma_1 \oplus \gamma_2$ , and  $\Gamma_3 = \gamma_1 \oplus \gamma_3$ . The contours are closed so that  $\text{Ind}_{\Gamma_k}(z)$  is defined for  $z = 0$ ,  $z = 3i/4$ , and  $z = -2$ .

The number  $\text{Ind}_\Gamma(z)$  is also called the *winding number* of the contour  $\Gamma$  around  $z$ . The above examples hint at the origin of the name.

### 3.2. Cauchy's theorem

**3.2.1 Cauchy's theorem for functions with primitives.** Suppose  $\Omega$  is a non-empty open set,  $f$  a continuous function on  $\Omega$  which has a primitive  $F$ . If  $\Gamma$  is a closed contour in  $\Omega$ , then  $\int_\Gamma f = 0$ .

**3.2.2 Cauchy's theorem for integer powers.** Suppose  $n \in \mathbb{Z}$ . Unless  $n = -1$  the power function  $z \mapsto z^n$  has the primitive  $z^{n+1}/(n+1)$  in either  $\Omega = \mathbb{C} \setminus \{0\}$  or  $\Omega = \mathbb{C}$  depending on whether  $n$  is negative or not. Consequently, if  $\Gamma$  is a closed contour in  $\Omega$  and if  $n \in \mathbb{Z} \setminus \{-1\}$ , then  $\int_\Gamma z^n dz = 0$ .

The exceptional case  $n = -1$  gives rise to interesting complications which we will discuss later. Recall, though, from 2.3.5 that  $\int_\gamma dz/z = 2\pi i$  when  $\gamma(t) = \exp(it)$  for  $t \in [0, 2\pi]$ .

**3.2.3 Triangles.** Let  $a, b, c$  be three pairwise distinct complex numbers. The set  $\Delta = \{ra + sb + tc : r, s, t \in [0, 1], r + s + t = 1\}$  is called a solid (possibly degenerate) *triangle* with vertices  $a$ ,  $b$ , and  $c$ . Solid triangles are closed subsets of  $\mathbb{C}$ . The line segments joining each pair of vertices are called *edges* of the triangle. They are obtained by setting  $r$ ,  $s$ , and  $t$  in

turn equal to zero and are thus subsets of  $\Delta$ . If  $c' = (a+b)/2$ ,  $b' = (a+c)/2$ , and  $a' = (b+c)/2$  (the midpoints of the edges), the sets  $\{a', b', c'\}$ ,  $\{a, b', c'\}$ ,  $\{a', b, c'\}$ , and  $\{a', b', c\}$  define respectively four new triangles  $\Delta_0, \dots, \Delta_3$ . Their union is  $\Delta$  and the intersection of any two of these is either a midpoint or one of the segments joining midpoints. Also  $\Delta_0, \dots, \Delta_3$  are **congruent** and are **similar** to  $\Delta$  itself. The **circumference** and the **diameter** of each one of the  $\Delta_k$  is half of that of  $\Delta$ . **Hint**

**3.2.4 Cauchy's theorem for triangles.** Suppose  $\Omega$  is a non-empty open set,  $z_0$  a point in  $\Omega$ , and  $f : \Omega \rightarrow \mathbb{C}$  a continuous function which is holomorphic on  $\Omega \setminus \{z_0\}$ . If  $\Delta$  is a solid triangle in  $\Omega$  with vertices  $a, b$ , and  $c$ , let  $\Gamma = \langle a, b, c, a \rangle$ . Then  $\int_{\Gamma} f = 0$ . **Hint**

**3.2.5 Cauchy's theorem for convex sets.** Suppose  $\Omega$  is a non-empty open convex set,  $z_0$  a point in  $\Omega$ , and  $f : \Omega \rightarrow \mathbb{C}$  a continuous function which is holomorphic on  $\Omega \setminus \{z_0\}$ . Fix  $a \in \Omega$  and define  $F : \Omega \rightarrow \mathbb{C}$  by  $F(z) = \int_{\langle a, z \rangle} f$ . Then  $F$  is an antiderivative of  $f$  and  $\int_{\Gamma} f = 0$  for every closed contour  $\Gamma$  in  $\Omega$ .

### 3.3. Consequences of Cauchy's theorem

**3.3.1 Cauchy's integral formula for convex sets.** Suppose  $\Omega$  is a non-empty open convex set and  $f : \Omega \rightarrow \mathbb{C}$  a holomorphic function on  $\Omega$ . If  $\Gamma$  is a closed contour in  $\Omega$  and  $z \in \Omega \setminus \Gamma^*$ , then

$$f(z) \operatorname{Ind}_{\Gamma}(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(u)}{u-z} du.$$

**3.3.2 Holomorphic functions are analytic.** We can now prove the central theorem of Complex Analysis.

**THEOREM.** A holomorphic function is analytic on its domain of definition.

From now on the words holomorphic and analytic may be considered synonymous (many authors do not ever make a distinction between them).

**3.3.3 Morera's<sup>1</sup> theorem.** Let  $\Omega$  be a non-empty open set. If  $f : \Omega \rightarrow \mathbb{C}$  is continuous and if  $\int_{\Gamma} f = 0$  for every triangular contour  $\Gamma \in \Omega$ , then  $f$  is holomorphic in  $\Omega$ .

**3.3.4 Sequences of holomorphic functions.** Let  $\Omega$  be a non-empty open set. Suppose  $n \mapsto f_n$  is a sequence of holomorphic functions defined on  $\Omega$  which converges uniformly to  $f : \Omega \rightarrow \mathbb{C}$ . Then  $f$  is holomorphic.

**3.3.5 General integral formulas for convex sets.** Suppose  $\Omega$  is a non-empty open convex set and  $f : \Omega \rightarrow \mathbb{C}$  a holomorphic function on  $\Omega$ . If  $\Gamma$  is a closed contour in  $\Omega$ ,  $z \in \Omega \setminus \Gamma^*$ , and  $n \in \mathbb{N}_0$ , then

$$f^{(n)}(z) \operatorname{Ind}_{\Gamma}(z) = \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(u)}{(u-z)^{n+1}} du.$$

**Hint**

**3.3.6 Cauchy's estimate.** If  $f : B(z_0, R) \rightarrow \mathbb{C}$  is a holomorphic function, then the radius of convergence of its Taylor series about  $z_0$  is at least equal to  $R$ . If  $|f(z)| \leq M$  whenever  $z \in B(z_0, R)$ , then

$$a_n = \frac{|f^{(n)}(z_0)|}{n!} \leq \frac{M}{R^n}$$

for all  $n \in \mathbb{N}_0$ .

---

<sup>1</sup>Giacinto Morera (1856 – 1909)

**3.3.7 Liouville's<sup>2</sup> theorem.** Every bounded entire function is constant.

**3.3.8 Zeros and their order.** Let  $z_0$  be a **zero** of the holomorphic function  $f$  and  $\sum_{n=0}^{\infty} a_n(z - z_0)^n$  the Taylor series of  $f$  about  $z_0$ . Then  $z_0$  is an isolated point in the set of all zeros of  $f$  unless  $a_n = 0$  for all  $n \in \mathbb{N}_0$ . If  $z_0$  is an isolated zero of  $f$ , the number  $m = \min(\{k \in \mathbb{N} : a_k \neq 0\})$  is called the *order* or the *multiplicity* of  $z_0$  as a zero of  $f$ . In this case we have  $f(z) = (z - z_0)^m g(z)$  where  $g$  is holomorphic in the domain of  $f$  and  $g(z_0) \neq 0$ .

**3.3.9 The set of zeros of a holomorphic function.** Let  $\Omega$  be an open connected subset of  $\mathbb{C}$  and  $f$  a holomorphic function on  $\Omega$ . Let  $Z(f)$  be the set of zeros of  $f$ , i.e.,  $Z(f) = \{a \in \Omega : f(a) = 0\}$ . Then either  $Z(f) = \Omega$  or else  $Z(f)$  is a set of isolated points.

**Hint**

**3.3.10 Analytic continuation.** Let  $f$  and  $g$  be holomorphic functions on  $\Omega$ , an open connected subset of  $\mathbb{C}$ . Then the following statements hold:

- (1) If  $f(z) = g(z)$  for all  $z$  in a set which contains a limit point of itself, then  $f = g$ .
- (2) If  $f^{(n)}(z_0) = g^{(n)}(z_0)$  for some  $z_0$  and all  $n \in \mathbb{N}_0$ , then  $f = g$ .

Suppose  $f$  is a holomorphic function on  $\Omega$  and that  $\Omega'$  is another open connected set which intersects  $\Omega$ . Then there is at most one holomorphic function on  $\Omega \cup \Omega'$  which coincides with  $f$  on  $\Omega$ . This function (if it exists) is called an *analytic continuation* of  $f$ .

### 3.4. The global version of Cauchy's theorem

**3.4.1 Continuity of parametric integrals.** Let  $\Omega$  be an open set,  $\Gamma$  a contour in  $\mathbb{C}$ , and  $\varphi$  a continuous complex-valued function on  $\Gamma^* \times \Omega$ . Define  $g(z) = \int_{\Gamma} \varphi(\cdot, z)$  on  $\Omega$ . Then  $g$  is continuous on  $\Omega$ .

**Hint**

**3.4.2 Differentiation under the integral.** Let  $\Omega$ ,  $\Gamma$ ,  $\varphi$ , and  $g$  be defined as in 3.4.1. Furthermore suppose that  $\varphi(u, \cdot)$  is holomorphic in  $\Omega$  for each  $u \in \Gamma^*$  and denote its derivative by  $\psi(u, \cdot)$ , so that  $\varphi(u, \cdot)$  is a primitive of  $\psi(u, \cdot)$  for every fixed  $u \in \Gamma^*$ . If  $\psi : \Gamma^* \times \Omega \rightarrow \mathbb{C}$  is continuous, then  $g$  is holomorphic in  $\Omega$ .

**Hint**

**3.4.3 A special (but important) case.** Suppose  $\Omega$  is an open subset of  $\mathbb{C}$ ,  $f$  is a holomorphic function on  $\Omega$ , and  $\Gamma$  is a contour in  $\Omega$ . Define  $\varphi, \psi : \Gamma^* \times \Omega \rightarrow \mathbb{C}$  by

$$\varphi(u, z) = \begin{cases} (f(u) - f(z))/(u - z) & \text{if } u \neq z, \\ f'(z) & \text{if } u = z \end{cases}$$

and

$$\psi(u, z) = \begin{cases} (f(u) - f(z) - f'(z)(u - z))/(u - z)^2 & \text{if } u \neq z, \\ \frac{1}{2}f''(z) & \text{if } u = z \end{cases}.$$

Then  $\varphi$  and  $\psi$  are both continuous on  $\Gamma^* \times \Omega$  and  $\psi(u, \cdot)$  is the derivative of  $\varphi(u, \cdot)$  whenever  $u \in \Gamma^*$ .

**Hint**

**3.4.4 Cauchy's integral formula, global version.** Suppose  $\Omega$  is an open subset of the complex plane,  $f$  is a holomorphic function on  $\Omega$ , and  $\Gamma$  is a closed contour in  $\Omega$  such that

---

<sup>2</sup>Joseph Liouville (1809 – 1882)

$\text{Ind}_\Gamma(w) = 0$  whenever  $w$  is not an element of  $\Omega$ . Then

$$f(z) \text{Ind}_\Gamma(z) = \frac{1}{2\pi i} \int_\Gamma \frac{f(u)}{u-z} du$$

for every  $z \in \Omega \setminus \Gamma^*$ .

**Hint**

**3.4.5 Cauchy's theorem, global version.** Suppose  $\Omega$ ,  $f$ , and  $\Gamma$  satisfy the same hypotheses as in 3.4.4. Then  $\int_\Gamma f = 0$ .

**3.4.6 Cauchy's theorem and integral formula for simply connected open sets.** A connected open set  $\Omega$  in the complex plane is called *simply connected* if  $\mathbb{C} \setminus \Omega$  has no bounded component. The set  $\mathbb{C} \setminus \{0\}$  is not simply connected but  $\mathbb{C} \setminus (-\infty, 0]$  is.

If  $\Omega$  is a non-empty open simply connected subset of  $\mathbb{C}$  and  $\Gamma$  a closed contour in  $\Omega$ , then  $\text{Ind}_\Gamma(w) = 0$  whenever  $w$  is not an element of  $\Omega$ . Consequently, if  $f$  is a holomorphic function on  $\Omega$ , then

$$f(z) \text{Ind}_\Gamma(z) = \frac{1}{2\pi i} \int_\Gamma \frac{f(u)}{u-z} du$$

for every  $z \in \Omega \setminus \Gamma^*$  and

$$\int_\Gamma f = 0.$$

**3.4.7 Laurent<sup>3</sup> series.** Suppose  $0 \leq r_1 < r_2$  and let  $f$  be holomorphic in the annulus  $\Omega = \{z \in \mathbb{C} : r_1 < |z-a| < r_2\}$ . Then,  $f$  can be expressed by a *Laurent series*, i.e.,

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z-a)^n.$$

This is actually an abbreviation for

$$f(z) = \sum_{n=0}^{\infty} a_n(z-a)^n + \sum_{n=1}^{\infty} a_{-n}(z-a)^{-n},$$

i.e., convergence of a Laurent series requires convergence of both series of positive and negative powers.

The coefficients  $a_n$  are given by the integrals

$$a_n = \frac{1}{2\pi i} \int_\gamma \frac{f(u)}{(u-a)^{n+1}} du$$

where  $\gamma = a + r' \exp(it)$ ,  $t \in [0, 2\pi]$  and  $r' \in (r_1, r_2)$ . **Hint**

The Taylor series of an analytic function is, of course, a special case of a Laurent series.

---

<sup>3</sup>Pierre Laurent (1813 – 1854)

## Isolated singularities

### 4.1. Classifying isolated singularities

Throughout this section  $\Omega$  denotes an open subset of the complex plane.

**4.1.1 Isolated singularities.** Suppose  $z_0 \in \Omega$  and  $f : \Omega \setminus \{z_0\} \rightarrow \mathbb{C}$  is holomorphic. Then  $z_0$  is called an *isolated singularity* of  $f$ .

The following are typical examples:  $z \mapsto (z^2 - z_0^2)/(z - z_0)$ ,  $z \mapsto 1/(z - z_0)$ , and  $z \mapsto \exp(1/(z - z_0))$ .

**4.1.2 Removable singularities.** Suppose  $z_0$  is an isolated singularity of a function  $f$ . If there is an analytic continuation of  $f$  to  $B(z_0, r)$  for some  $r > 0$ , then  $z_0$  is called a *removable singularity* of  $f$ .

Suppose  $z_0$  is an isolated singularity of  $f$ . It is a removable singularity of  $f$  if and only if  $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$ . **Hint**

**4.1.3 Poles.** Suppose  $z_0$  is an isolated singularity of a function  $f$ . If  $\lim_{z \rightarrow z_0} 1/f(z) = 0$  (to avoid saying  $\lim_{z \rightarrow z_0} f(z) = \infty$ ), then  $z_0$  is called a *pole* of  $f$ .

Suppose  $z_0$  is an isolated singularity of  $f$ . It is a pole of  $f$  if and only if there is a natural number  $m$  such that  $z_0$  is a removable singularity of  $z \mapsto (z - z_0)^m f(z)$ . The smallest such  $m$  is called the *order* or the *multiplicity* of  $z_0$  as a pole of  $f$ .

Suppose  $f$  is holomorphic on  $B(z_0, r) \setminus \{z_0\}$ .  $f$  has a pole of order  $m$  at  $z_0$  if and only if  $1/f$  (properly extended to  $B(z_0, r)$ ) has a zero of order  $m$  at  $z_0$ .

**4.1.4 Essential singularities.** An isolated singularity of a function which is neither removable nor a pole is called an *essential singularity* of the function.

**4.1.5 Meromorphic functions.** If  $P$  is a set of isolated points, if  $f$  is holomorphic on  $\Omega \setminus P$ , and if no point of  $P$  is an essential singularity of  $f$ , then  $f$  is called *meromorphic* on  $\Omega$ .

**4.1.6 The Casorati<sup>1</sup>-Weierstrass theorem.** If  $z_0 \in \Omega$  is an essential singularity of  $f : \Omega \setminus \{z_0\} \rightarrow \mathbb{C}$  and  $r > 0$  is such that  $B' = B(z_0, r) \setminus \{z_0\} \subset \Omega$ , then  $f(B')$  is **dense** in  $\mathbb{C}$ . **Hint**

**4.1.7 Laurent series about an isolated singularity.** A **punctured disk**  $\{z \in \mathbb{C} : 0 < |z - z_0| < R\}$  is a special case of an annulus and thus a holomorphic function defined on it has a Laurent series expansion. In particular, if  $z_0$  is an isolated singularity of  $f$  then there is a punctured disk  $\{z \in \mathbb{C} : 0 < |z - z_0| < R\}$  (possibly the punctured plane) on which

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n.$$

---

<sup>1</sup>Felice Casorati (1835 – 1890)

The point  $z_0$  is a removable singularity if and only if  $a_n = 0$  for all  $n < 0$ . It is a pole of order  $m$  if and only if  $a_n = 0$  for all  $n < -m$  and  $a_{-m} \neq 0$ . It is an essential singularity if and only if the set  $\{n \in \mathbb{Z} : a_n \neq 0\}$  is not bounded below.

If  $f$  has a pole of order  $m$  at  $z_0$ , then there are complex numbers  $c_1, \dots, c_m$  so that

$$z \mapsto f(z) - \sum_{k=1}^m \frac{c_k}{(z - z_0)^k}$$

has a removable singularity at  $z_0$ . The numbers  $c_1, \dots, c_m$  are unique and the sum  $\sum_{k=1}^m c_k/(z - z_0)^k$  is called the *principal part* or *singular part* of  $f$  at  $z_0$ .

## 4.2. The calculus of residues

**4.2.1 Residues.** If the holomorphic function  $f$  has the isolated singularity  $z_0$  and the Laurent expansion  $\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ , then the number

$$a_{-1} = \frac{1}{2\pi i} \int_{\gamma} f,$$

where  $\gamma$  traces a sufficiently small circle about  $z_0$ , is called the *residue* of  $f$  at  $z_0$  and is denoted by  $\text{Res}(f, z_0)$ .

If  $\lim_{z \rightarrow z_0} (z - z_0)f(z) = a \neq 0$ , then  $z_0$  is a *simple pole*, i.e., a pole of order one, of  $f$  and  $a = \text{Res}(f, z_0)$ .

**4.2.2 The residue theorem.** Suppose  $\Omega$  is an open subset of the complex plane,  $f$  is a holomorphic function on  $\Omega' = \Omega \setminus \{z_1, \dots, z_n\}$ , and  $\Gamma$  is a closed contour in  $\Omega'$  such that  $\text{Ind}_{\Gamma}(w) = 0$  whenever  $w$  is not an element of  $\Omega$  (i.e.,  $\Gamma$  does not wind around any point outside  $\Omega$ ). Then

$$\sum_{k=1}^n \text{Res}(f, z_k) \text{Ind}_{\Gamma}(z_k) = \frac{1}{2\pi i} \int_{\Gamma} f.$$

Hint

### 4.2.3 Some examples.

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{1}{1+x^4} dx \\ & \int_{-\infty}^{\infty} \frac{1}{\exp(x) + \exp(-x)} dx \\ & \int_0^{2\pi} \frac{1}{2 + \sin x} dx \end{aligned}$$

Hints

**4.2.4 Counting zeros and poles.** Suppose  $f$  is a non-constant meromorphic function on the open and connected set  $\Omega$ ,  $Z$  is the set of its zeros, and  $P$  is the set of its poles. For any point  $z_0 \in \Omega$  there is a unique integer  $M(z_0)$  denoting the smallest index for which the coefficient in the Laurent expansion of  $f$  about  $z_0$  is non-zero. In fact, if  $z_0$  is a zero or a pole of  $f$  its multiplicity is  $M(z_0)$  or  $-M(z_0)$ , respectively.

Let  $\Gamma$  be a closed contour in  $\Omega \setminus (Z \cup P)$  such that  $\text{Ind}_{\Gamma}(w) = 0$  whenever  $w \in \mathbb{C} \setminus \Omega$ . Then

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'}{f} = \sum_{z \in Z \cup P} M(z) \text{Ind}_{\Gamma}(z).$$

In applying this result note that there are at most finitely many zeros and poles in those components of  $\Omega$  which have a non-zero index, so that the sums occurring are actually finite sums.

In particular, if  $f$  is holomorphic on  $\Omega$  and  $\text{Ind}_\Gamma$  assumes only the values 0 and 1, then  $\frac{1}{2\pi i} \int_\Gamma \frac{f'}{f}$  is the number of zeros (with multiplicities) in the set  $\{z \in \mathbb{C} : \text{Ind}_\Gamma(z) = 1\}$ .

**4.2.5 Rouché's<sup>2</sup> theorem.** Let  $f$  and  $g$  be meromorphic functions on the open set  $\Omega$  and assume that  $\overline{B}(a, r) \subset \Omega$ . Let  $n_z(f)$  be the number of zeros of  $f$  in  $B(a, r)$ , counted according to their multiplicity. Similarly,  $n_p(f)$  is the count of poles, and  $n_z(g)$  and  $n_p(g)$  are the analogous quantities for  $g$ . If no zero or pole lies on the circle  $C = \{z : |z - a| = r\}$  and if  $|f(z) - g(z)| < |f(z)| + |g(z)|$  for all  $z \in C$ , then  $n_z(f) - n_p(f) = n_z(g) - n_p(g)$ . **Hint**

**4.2.6 The open mapping theorem.** Suppose  $f$  is a non-constant holomorphic function on the open and connected set  $\Omega$ . Then  $f(\Omega')$  is open whenever  $\Omega'$  is an open subset of  $\Omega$ . **Hint**

**4.2.7 The maximum modulus theorem.** Suppose  $f$  is a holomorphic function on  $\Omega$  and  $|f(z_0)|$  is a **local maximum** of  $|f|$ . Then  $f$  is constant.

---

<sup>2</sup>Eugène Rouché (1832 – 1910)





## CHAPTER 5

### A zoo of functions

In this chapter we investigate briefly the most elementary functions of analysis. Polynomials and exponential functions have been introduced earlier since they are too important to postpone their use.

#### 5.1. Polynomial functions

**5.1.1 Polynomial functions.** If  $n$  is a non-negative integer and  $a_0, a_1, \dots, a_n$  are complex numbers, then the function  $p : \mathbb{C} \rightarrow \mathbb{C}$  defined by

$$p(z) = \sum_{k=0}^n a_k z^k$$

is called a *polynomial* (of a single variable). The integer  $n$  is called the *degree* of  $p$  if  $a_n$ , the leading coefficient of  $p$ , is different from 0. The zero function is also a polynomial but no degree is assigned to it. Any polynomial is an entire function.

If  $p$  and  $q$  are polynomials of degree  $n$  and  $k$ , respectively, then  $p + q$  and  $pq$  are also polynomials. The degree of  $p + q$  is the larger of the numbers  $n$  and  $k$  unless  $n = k$  in which case the degree is at most  $n$ . The degree of  $pq$  equals  $n + k$ .

**5.1.2 The fundamental theorem of algebra.** Suppose  $p$  is a polynomial of degree  $n \geq 1$ . Then there exist numbers  $a$  and  $z_1, \dots, z_n$  (not necessarily distinct) such that

$$p(z) = a \prod_{k=1}^n (z - z_k).$$

**Hint**

**5.1.3 Zeros of polynomials.** The coefficients  $a_k$  of a **monic** polynomial are given as **symmetric polynomials** in terms of the zeros, e.g.,  $a_{n-1} = -\sum_{k=1}^n z_k$  and  $a_0 = (-1)^n \prod_{k=1}^n z_k$ . They vary continuously with the zeros. **Hint**

That the converse is also true can be proved with the aid of Rouché's theorem. The precise statement is as follows: If  $z_0$  is a zero of order  $m$  of the polynomial  $f(z) = \sum_{k=0}^n a_k z^k$  and if  $f$  has no other zeros in  $\overline{B}(z_0, r)$  then the following holds: For every  $\varepsilon \in (0, r)$  there is a  $\delta > 0$  such that, if  $|a_k - b_k| < \delta$  for  $k = 0, \dots, n$  and  $g(z) = \sum_{k=0}^n b_k z^k$ , then  $g$  has precisely  $m$  zeros in  $B(z_0, \varepsilon)$  (counting multiplicities).

**5.1.4 Among the entire functions only polynomials grow like powers.** Suppose  $f$  is an entire function and  $k \mapsto r_k$  is an increasing unbounded sequence of positive numbers. Furthermore, suppose there are numbers  $N \in \mathbb{N}$  and  $C > 0$  such that  $|f(z)| \leq C|z|^N$  for all  $z$  lying on any of the circles  $|z| = r_k$ . Then  $f$  is a polynomial of degree at most  $N$ .

It is in fact enough to require  $\operatorname{Re}(f(z)) \leq C|z|^N$  to arrive at the same conclusion. **Hint**

Thus the real (or the imaginary part) of a polynomial grows roughly at the same pace as the modulus. This is, in fact, a special case of a more general result, the Borel-Carathéodory<sup>1</sup> theorem, which allows to estimate the modulus of an entire function by its real (or imaginary) part.

## 5.2. Rational functions

**5.2.1 The Riemann sphere.** The set  $S^2 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1\}$  is called the *unit sphere*. The point  $(0, 0, 1)$  is called the *north pole* of the sphere. Let  $(x, y, z)$  be a point in  $S^2$  other than the north pole. The map  $t \mapsto (tx, ty, tz + 1 - t)$ ,  $t \in \mathbb{R}$ , describes a straight line passing through the north pole and the point  $(x, y, z)$ . This line intersects the plane  $z = 0$  in the unique point  $(x/(1 - z), y/(1 - z), 0)$ . We have thus defined the map

$$\pi : S^2 \setminus \{(0, 0, 1)\} \rightarrow \mathbb{C} : (x, y, z) \mapsto \frac{x + iy}{1 - z}.$$

This map, called the *stereographic projection*, is bijective. Points in the southern hemisphere correspond to points in the **unit disk**, points on the equator correspond to points on the **unit circle**, and points in the northern hemisphere correspond to points outside the closed unit disk.

With this interpretation the sphere  $S^2$  is called the *Riemann sphere* and is denoted by  $\mathbb{C}_\infty$ . Since  $S^2$  is a compact metric space (as a subspace of  $\mathbb{R}^3$ ),  $\mathbb{C}_\infty$  is a one-point compactification of  $\mathbb{C}$ . Note that points on the sphere close to the north pole are those whose image points in the plane have large absolute values.

We can now view a meromorphic function on  $\Omega$  as a function from  $\Omega$  to  $\mathbb{C}_\infty$ . As such it is continuous at every point of  $\Omega$  even the poles.

**5.2.2 Rational functions.** Let  $p$  and  $q$  be polynomials and  $Q = \{z \in \mathbb{C} : q(z) = 0\}$ . Assume  $Q \neq \mathbb{C}$ , i.e.,  $q$  is not the zero polynomial. Then, the function  $r : (\mathbb{C} \setminus Q) \rightarrow \mathbb{C}$  given by

$$r(z) = \frac{p(z)}{q(z)}$$

is called a *rational function*.

A rational function may also be interpreted as a function from  $\mathbb{C}_\infty$  to  $\mathbb{C}_\infty$  by setting  $r(z) = \infty$  if  $z \in Q$  and  $r(\infty) = \lim_{z \rightarrow \infty} r(z)$  (which may again be infinity).

**5.2.3 Möbius transforms.** A rational function of the type  $z \mapsto (az + b)/(cz + d)$  where  $ad - bc \neq 0$  is called a *Möbius transform*<sup>2</sup>. A Möbius transform may be interpreted as a bijective map from the Riemann sphere to itself. The set of all Möbius transforms forms a **group** under composition. Special cases of Möbius transforms are the translations  $z \mapsto z + b$ , the dilations and rotations  $z \mapsto az$  where  $a > 0$  and  $|a| = 1$ , respectively, and the inversion  $z \mapsto 1/z$ . Any Möbius transform is a composition of at most five of these.

Möbius transforms have many interesting properties to which an entire chapter might be devoted.

## 5.3. Exponential and trigonometric functions

**5.3.1 The exponential function.** We defined the exponential function already in **2.5.6**. Hence most of the following will be a repetition of results obtained there. The function

<sup>1</sup>Constantin Carathéodory (1873 – 1950)

<sup>2</sup>August Ferdinand Möbius (1790 – 1868)

$\exp : \mathbb{C} \rightarrow \mathbb{C}$  defined by

$$\exp(z) = \sum_{k=0}^{\infty} \frac{z^k}{k!}$$

is called the *exponential function*. It is an entire function and  $\exp'(z) = \exp(z)$ .

If  $x, y \in \mathbb{C}$ , then  $\exp(x + y) = \exp(x)\exp(y)$ . In particular,  $\exp(z)$  is never zero and  $\exp(-z) = 1/\exp(z)$ .

The exponential function has period  $2\pi i$ , i.e.,  $\exp(z + 2\pi i) = \exp(z)$ . If  $p$  is any period of  $\exp$ , then there is an integer  $m$  such that  $p = 2m\pi i$ . Let  $a$  be a fixed real number and  $S_a = \{z \in \mathbb{C} : a < \operatorname{Im}(z) \leq a + 2\pi\}$ . Then  $\exp|_{S_a}$  maps  $S_a$  bijectively to  $\mathbb{C} \setminus \{0\}$ .

**5.3.2 Trigonometric functions.** The trigonometric functions are defined by

$$\cos(z) = \frac{\exp(iz) + \exp(-iz)}{2},$$

and

$$\sin(z) = \frac{\exp(iz) - \exp(-iz)}{2i}.$$

They are entire functions, called the *cosine* and *sine function*, respectively.

The following properties, familiar from the real case, extend to the complex case:

- (1) Derivatives:  $\sin'(z) = \cos(z)$  and  $\cos'(z) = -\sin(z)$ .
- (2) The Pythagorean theorem:  $(\sin z)^2 + (\cos z)^2 = 1$  for all  $z \in \mathbb{C}$ .
- (3) Addition theorems:

$$\sin(z + w) = \sin(z)\cos(w) + \cos(z)\sin(w)$$

and

$$\cos(z + w) = \cos(z)\cos(w) - \sin(z)\sin(w)$$

for all  $z, w \in \mathbb{C}$ . (4) Taylor series:

$$\sin(z) = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n+1}}{(2n+1)!}$$

and

$$\cos(z) = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{(2n)!}.$$

In particular,  $\sin$  and  $\cos$  are extensions of the functions defined in Real Analysis.

- (5)  $\cos(z) = \sin(z + \pi/2)$ .
- (6)  $\sin(z) = 0$  if and only if  $z$  is an integer multiple of  $\pi$ .
- (7) The range of both, the sine and the cosine function, is  $\mathbb{C}$ .

## 5.4. The logarithmic function and powers

**5.4.1 The real logarithm.** Since  $\exp : \mathbb{R} \rightarrow (0, \infty)$  is a bijective function, it has an inverse  $\ln : (0, \infty) \rightarrow \mathbb{R}$ . One shows that  $\ln(1) = 0$  and  $\ln'(x) = 1/x$  so that, by the fundamental theorem of calculus

$$\ln(x) = \int_1^x \frac{dt}{t}$$

for all  $x > 0$ .

**5.4.2 The logarithm.** Suppose  $\Omega$  is a simply connected non-empty open subset of  $\mathbb{C} \setminus \{0\}$  and  $\Gamma$  is a contour in  $\mathbb{C} \setminus \{0\}$  connecting 1 to  $z_0 \in \Omega$ . Then we define the function  $L_\Gamma : \Omega \rightarrow \mathbb{C}$  by setting

$$L_\Gamma(z) = \int_{\Gamma \oplus \beta} \frac{du}{u}$$

where  $\beta$  is a contour in  $\Omega$  connecting  $z_0$  to  $z$ . As the notation suggests  $L_\Gamma(z)$  does not depend on  $\beta$  as long as  $\beta$  remains in  $\Omega$ . The function  $L_\Gamma$  is called a *branch of the logarithm* on  $\Omega$ .

Suppose now that  $\Gamma'$  is another contour in  $\mathbb{C} \setminus \{0\}$  connecting 1 to a point in  $\Omega$ . If  $z'_0$  is the end point of  $\Gamma'$  and  $\gamma$  is a contour in  $\Omega$  connecting  $z_0$  and  $z'_0$ , then

$$L_\Gamma(z) - L_{\Gamma'}(z) = 2\pi i \operatorname{Ind}_{\Gamma \oplus \gamma \ominus \Gamma'}(0).$$

Thus there are (at most) countably many different functions  $L_\Gamma$  which are defined this way, even though there are many more contours  $\Gamma$  connecting 1 to a point in  $\Omega$ . In fact, for each  $m \in \mathbb{Z}$  there is a  $\Gamma'$  such that  $L_\Gamma(z) - L_{\Gamma'}(z) = 2m\pi i$ . This shows that defining the logarithm as an antiderivative comes with certain difficulties (to say the least). At the same time this behavior gives rise to a lot of interesting mathematics.

$L_\Gamma$  is a holomorphic function on  $\Omega$  with derivative  $1/z$ . Moreover,  $\exp(L_\Gamma(z)) = z$  for all  $z \in \Omega$  but  $L_\Gamma(\exp(z))$  may well differ from  $z$  by an integer multiple of  $2\pi i$ .

**5.4.3 The principal branch of the logarithm.** Let  $\Omega = \mathbb{C} \setminus (-\infty, 0]$ . Then

$$\log(z) = \int_\gamma \frac{du}{u},$$

where  $\gamma$  is contour in  $\Omega$  connecting 1 and  $z$ , is uniquely defined for any  $z \in \Omega$  (i.e.,  $\log = L_\Gamma$  where  $\Gamma : [0, 1] \rightarrow \mathbb{C} : t \mapsto 1$ ). It is called the principal branch of the *logarithm*. If  $z \in \Omega$  has polar representation  $z = r \exp(it)$  where  $t \in (-\pi, \pi)$  and  $r > 0$ , then

$$\log(z) = \ln(r) + it.$$

The range of  $\log$  is the strip  $\{z \in \mathbb{C} : |\operatorname{Im}(z)| < \pi\}$ . In particular,  $\log(\exp(z)) = z$  if and only if  $|\operatorname{Im}(z)| < \pi$ .

The Taylor series of  $z \mapsto \log(1+z)$  about  $z = 0$  has radius of convergence 1 and is given by

$$\log(1+z) = -\sum_{n=1}^{\infty} \frac{(-z)^n}{n}.$$

If  $\operatorname{Re}(a)$  and  $\operatorname{Re}(b)$  are positive, then  $\log(ab) = \log(a) + \log(b)$ . The conclusion may be wrong when the hypothesis is not satisfied.

**5.4.4 Powers.** If  $L : \Omega \rightarrow \mathbb{C}$  is a branch of the logarithm, if  $a \in \Omega$  and  $b \in \mathbb{Z}$ , then  $a^b = \exp(bL(a))$ . We may therefore define

$$a^b = \exp(bL(a))$$

for any  $b \in \mathbb{C}$ . Be aware, though, that the value of  $a^b$  depends on the branch of the logarithm chosen and is therefore ambiguous when  $b \notin \mathbb{Z}$ . For example,  $z^{1/2}$  may represent two different numbers. In most cases one chooses, of course, the principal branch of the logarithm and defines  $a^b = \exp(b \log(a))$  (forsaking the definition of powers of negative numbers). In this case we get  $e^z = \exp(z)$  for all  $z \in \mathbb{C}$ , after we define  $e = \exp(1)$ .

**5.4.5 Power functions.** Let  $p$  be a complex number and  $L : \Omega \rightarrow \mathbb{C}$  a branch of the logarithm. The function  $\Omega \rightarrow \mathbb{C} : z \mapsto z^p = \exp(pL(z))$  is called a *branch of the power function*. Each such branch is a holomorphic function which never vanishes. Note that  $1/z^p = z^{-p}$ .

If  $p \in \mathbb{N}_0$  all branches of the associated power function may be analytically extended to the entire complex plane and any branch gives then rise to one and the same function. The same is true if  $p \in \mathbb{Z}$  is negative, except that the function may not be extended to 0 since 0 is then a pole of the function.

If  $p \in \mathbb{Q}$  there are only finitely many different branches of  $z \mapsto z^p$  in any simply connected open set. In fact, if  $m/n$  is a representation of  $p$  in lowest terms, there will be  $n$  branches of  $z \mapsto z^p$ .

**5.4.6 Holomorphic functions without zeros.** Suppose  $f$  is a holomorphic function defined on a simply connected open set  $\Omega$ . If  $f$  has no zeros, there is an holomorphic function  $g$  on  $\Omega$  such that  $f(z) = \exp(g(z))$ . **Hint**

**5.4.7 A definite integral.** We will later use the following result (which is also an instructive example):

$$\int_{-\pi}^{\pi} \log |1 - \exp(it)| dt = 0.$$

**Hint**



## CHAPTER 6

# Entire functions

### 6.1. Infinite products

**6.1.1 Infinite products.** If  $n \mapsto z_n$  is a sequence of complex numbers, we denote the sequence  $k \mapsto p_k = \prod_{n=1}^k z_n$  of partial products by  $\prod_{n=1}^{\infty} z_n$ . If  $\lim_{k \rightarrow \infty} p_k$  exists, we will call it the *infinite product* of the numbers  $z_n$  and also denote it by  $\prod_{n=1}^{\infty} z_n$  (abusing notation as we do for series).

**6.1.2 Convergence criteria for infinite products.** Suppose  $\sum_{n=1}^{\infty} (z_n - 1)$  is absolutely convergent. Then there is an  $n_0 \in \mathbb{N}$  such that  $|z_n - 1| < 1/2$  for all  $n \geq n_0$ . Moreover,  $\sum_{n=n_0}^{\infty} \log(z_n)$  is absolutely convergent and  $\prod_{n=n_0}^{\infty} z_n$  converges to a non-zero number. The latter limit is independent of the order of the factors since

$$\prod_{n=n_0}^{\infty} z_n = \exp\left(\sum_{n=n_0}^{\infty} \log(z_n)\right).$$

We will then say that  $\prod_{n=1}^{\infty} z_n$  *converges absolutely*.

If  $\prod_{n=1}^{\infty} z_n$  converges absolutely to 0, then finitely many and only finitely many of the  $z_n$  vanish.

**6.1.3 Infinite products of analytic functions.** Let  $\Omega$  be a non-empty open set and  $n \mapsto f_n$  a sequence of holomorphic functions on  $\Omega$  none of which is identically equal to zero. If  $\sum_{n=1}^{\infty} |f_n - 1|$  converges uniformly on  $\Omega$ , then  $\prod_{n=1}^{\infty} f_n$  converges to a holomorphic function  $f$  on  $\Omega$ . If  $z_0$  is a zero of  $f$ , then it is a zero of only finitely many of the functions  $f_n$  and the order of  $z_0$  as a zero of  $f$  is the sum of the orders of  $z_0$  as a zero of  $f_n$ .

### 6.2. Weierstrass's factorization theorem

**6.2.1 Elementary factors.** The functions

$$\begin{aligned} E_0(z) &= 1 - z \\ E_p(z) &= (1 - z) \exp(z + \dots + z^p/p) \end{aligned}$$

are called *elementary factors*. Note that  $E_p(\cdot/a)$  is entire and has a simple zero at  $a$  but no other zero.

**6.2.2 Estimates for elementary factors.** If  $|z| \leq 1/2$ , then  $|E_p(z) - 1| \leq 2e|z|^{p+1}$  and

$$-2e|z|^{p+1} \leq \log(|E_p(z)|) \leq 2e|z|^{p+1}.$$

Also, if  $|z| \geq 1/2$ , then

$$-c|z|^p \leq \log(|E_p(z)/(1 - z)|) \leq c|z|^p$$

for some  $c > 0$ .

**6.2.3 Prescribing zeros of an entire function.** Suppose  $a_n$  is a (finite or infinite) sequence of non-zero complex numbers with no finite limit point. Then there is a sequence  $n \mapsto p_n \in \mathbb{N}_0$ , e.g.,  $p_n = n$ , such that

$$\forall r > 0 : \sum_n (r/|a_n|)^{p_n+1} < \infty.$$

Moreover,

$$\prod_n E_{p_n}(z/a_n)$$

converges to an entire function  $f$ . The zeros of  $f$  are precisely the numbers  $a_n$  and the order of  $z_0$  as a zero of  $f$  equals  $\#\{n : z_0 = a_n\}$ .

**6.2.4 The Weierstrass factorization theorem.** Let  $f$  be an entire function and suppose that  $a_n$ ,  $n \in \mathbb{N}$ , are the non-zero zeros of  $f$  repeated according to their multiplicity. Then there is an integer  $m$ , an entire function  $g$ , and a sequence  $n \mapsto p_n \in \mathbb{N}_0$  such that

$$f(z) = z^m \exp(g(z)) \prod_n E_{p_n}(z/a_n).$$

### 6.3. Counting zeros using Jensen's theorem

**6.3.1 Counting zeros.** If  $f$  is an entire function we define  $n_f(r)$  to be the number of zeros, counting multiplicities, of  $f$  in  $\overline{B}(0, r)$ .

**6.3.2 Jensen's<sup>1</sup> theorem.** Suppose  $f$  is an entire function and  $f(0) \neq 0$ . Denote its non-zero zeros, repeated according to their multiplicity and ordered by size, by  $a_k$ ,  $k \in \mathbb{N}$ . Then

$$\sum_{k=1}^{n(r)} \log(r/|a_k|) + \log |f(0)| = \frac{1}{2\pi} \int_0^{2\pi} \log |f(r \exp(it))| dt.$$

**6.3.3 Order of an entire function.** An entire function is said to be of finite order if the set

$$S = \{a \in [0, \infty) : \exists r_0 > 0 : |z| > r_0 \implies |f(z)| \leq \exp(|z|^a)\}$$

is not empty. The number  $\inf S$  is called the *order* of  $f$ .

**6.3.4 The growth of an entire function controls the number of its zeros.** Let  $f$  be an entire function of finite order  $\rho$  for which  $f(0) = 1$  and let  $\varepsilon$  be a positive number. Then there is an  $R > 0$  such that

$$n_f(r) \leq (er)^{\rho+\varepsilon}$$

for all  $r \geq R$ .

Hence if  $s > \rho$  we may choose  $\varepsilon \in (0, s - \rho)$  and obtain

$$\frac{1}{r^s} \leq \frac{e^s}{n_f(r)^{s/(\rho+\varepsilon)}}.$$

From this it follows that  $\sum_{n \in \mathbb{N}} |a_n|^{-s} < \infty$  whenever  $s > \rho$ .

---

<sup>1</sup>Johan Jensen (1859 – 1925)



**6.3.5 Exponent of convergence.** Let  $n \mapsto a_n$  be a sequence of non-zero complex numbers with no finite limit point. Then

$$\tau = \inf\{s > 0 : \sum_{n=1}^{\infty} |a_n|^{-s} < \infty\},$$

if it exists, is called the *exponent of convergence* of the sequence  $n \mapsto a_n$ .

If  $f$  is an entire function of finite order  $\rho$  and if the exponent of convergence of the non-zero zeros of  $f$  is  $\tau$ , then  $\tau \leq \rho$ . In this case Weierstrass's factorization theorem takes the form

$$f(z) = z^m \exp(g(z)) \prod_n E_p(z/a_n)$$

where  $p \in \mathbb{N}_0$  satisfies  $p \leq \tau < p + 1$ .

#### 6.4. Hadamard's factorization theorem

**6.4.1 Canonical products.** Let  $n \mapsto a_n$  be a sequence of non-zero complex numbers without a finite limit point whose exponent of convergence,  $\tau$ , is finite. If  $p \in \mathbb{N}_0$  is such that  $p \leq \tau < p + 1$  then  $g(z) = \prod_n E_p(z/a_n)$  is called the *canonical product* for the sequence  $n \mapsto a_n$ .

The function  $g$  is an entire function of order  $\tau$ .

**6.4.2 Controlling decay of canonical products.** Suppose  $a_n$  is a (finite or infinite) sequence of non-zero complex numbers and denote its exponent of convergence by  $\tau$ . Let  $p \in \mathbb{N}_0$  be such that  $p \leq \tau < p + 1$ . If  $\tau < s < p + 1$  and  $|z - a_n| \geq |a_n|^{-p-1}$  for all  $n \in N$ , then

$$\prod_n |E_p(z/a_n)| \geq \exp(-c|z|^s)$$

for some  $c > 0$ .

Between any two large natural numbers  $k$  and  $k + 1$  there is a number  $r_k$  such that  $|z - a_n| \geq |a_n|^{-p-1}$  for all  $z$  on the circle  $|z| = r_k$  and all  $n \in \mathbb{N}$ .

**6.4.3 Hadamard's Factorization Theorem.** Let  $f$  be an entire function of order  $\rho$  and zeros by  $a_n$ ,  $n \in N$ . Assume that the  $a_n$  are repeated according to their multiplicities and are arranged such that  $|a_1| \leq |a_2| \leq \dots$ . Let  $p = \lfloor \rho \rfloor$ , the largest integer not exceeding  $\rho$  and  $m \in \mathbb{N}_0$  the order of 0 as a zero of  $f$ . Then there exists a polynomial  $g$  of degree at most  $p$  such that

$$f(z) = z^m \exp(g(z)) \prod_{n \in N} E_p(z/a_n).$$

**6.4.4 The very little Picard theorem.** An entire function of finite order that misses two values is constant.

Picard's<sup>2</sup> little theorem states that this so for every entire function. Picard's big theorem states that near an isolated singularity a holomorphic function takes every value, with at most one exception, infinitely often; this is an extension of the Casorati-Weierstrass theorem.

**6.4.5 An entire function of non-integral order assumes every value infinitely often.**

---

<sup>2</sup>Émile Picard (1856 – 1941)



## APPENDIX A

### Topology in metric spaces

The following material is covered in a course on topology.

#### A.1. Basics

**A.1.1 Topology for a set.** Let  $X$  be a set. The system  $\tau$  of subsets of  $X$  is called a *topology* for  $X$  if it has the following properties:

- (1)  $\emptyset$  and  $X$  are in  $\tau$
- (2) Any union of elements of  $\tau$  is again in  $\tau$ .
- (3) The intersection of two elements of  $\tau$  is again in  $\tau$ .

If  $\tau$  is a topology for  $X$ , the pair  $(X, \tau)$  is called a *topological space*. The elements of  $\tau$  are called *open sets*, while their complements are called *closed sets*.

**A.1.2 Metric spaces.** Let  $X$  be a set. A function  $d : X \times X \rightarrow [0, \infty)$  is called a *metric* or a *distance function* in  $X$  if it has the following properties:

- (1)  $d(x_1, x_2) = 0$  if and only if  $x_1 = x_2$ .
- (2)  $d(x_1, x_2) = d(x_2, x_1)$  for all  $x_1, x_2 \in X$ .
- (3)  $d(x_1, x_3) \leq d(x_1, x_2) + d(x_2, x_3)$  for all  $x_1, x_2, x_3 \in X$  (*triangle inequality*).

$(X, d)$  is then called a *metric space* and the number  $d(x_1, x_2)$  is called the *distance* between  $x_1$  and  $x_2$ .

If  $X'$  is a subset of  $X$  then  $(X', d')$  is again a metric space when  $d'$  is the restriction of  $d$  to  $X' \times X'$ .

If  $\mathbb{R}^n$  is equipped with the distance function  $d$  defined by  $d(x, y)^2 = \sum_{k=1}^n (x_k - y_k)^2$ , it becomes a metric space called the Euclidean space.

**A.1.3 Open and closed balls.** Let  $(X, d)$  be a metric space,  $x_0$  a point in  $X$ , and  $r$  a non-negative real number. Then the set  $\underline{B}(x_0, r) = \{x \in X : d(x, x_0) < r\}$  is called the open *ball* of radius  $r$  centered at  $x_0$  while  $\overline{B}(x_0, r) = \{x \in X : d(x, x_0) \leq r\}$  is called the closed ball of radius  $r$  centered at  $x_0$ . Note that  $\underline{B}(x_0, 0) = \emptyset$  so that the empty set is an open ball. Similarly, the *singleton*  $\{x_0\} = \overline{B}(x_0, 0)$  is a closed ball.

A subset of a metric space is called *bounded* if it is contained in some ball.

**A.1.4 Metric spaces are topological spaces.** Let  $(X, d)$  be a metric space. Then

$$\tau = \{U \subset X : U \text{ is a union of open balls}\}$$

is a topology for  $X$ . This follows easily once one proves with the help of the triangle inequality that the intersection of two open balls is a union of open balls and hence an open set.

**A.1.5 Sequences and their limits.** Let  $(X, d)$  be a metric space. A function  $x : \mathbb{N} \rightarrow X : n \mapsto x_n$  is called a *sequence* in  $X$ . We say that the sequence  $x$  is *convergent* if the

following statement holds:

$$\exists L \in X : \forall \varepsilon > 0 : \exists N > 0 : \forall n > N : d(x_n, L) < \varepsilon.$$

If a sequence is not convergent, we say that it *diverges* or is *divergent*.

If a sequence  $x$  is convergent there is only one element in  $X$  to which it converges. This element is called the *limit* of the sequence  $x$  and is denoted by  $\lim_{n \rightarrow \infty} x_n$ .

**A.1.6 Continuity.** Let  $(X, d)$  and  $(Y, \rho)$  be metric spaces and  $f$  a function from  $X$  to  $Y$ . Then  $f$  is called *continuous* at  $a \in X$  if the following statement is true:

$$\forall \varepsilon > 0 : \exists \delta > 0 : \forall x \in X : d(x, a) < \delta \Rightarrow \rho(f(x), f(a)) < \varepsilon.$$

$f$  is called *continuous* if it is continuous at every point of  $X$ .

It follows that  $f$  is continuous if and only if the pre-image of any open subset of  $Y$  is an open subset of  $X$ .

## A.2. Compactness

**A.2.1 Compact sets.** A subset  $S$  of a topological space is called *compact* if every *open cover* of  $S$  has a *finite subcover*. More precisely, whenever  $\mathcal{V}$  is a family of open sets such that  $S \subset \bigcup_{V \in \mathcal{V}} V$  then there is a finite subset  $\{V_1, \dots, V_n\}$  of  $\mathcal{V}$  such that  $S \subset \bigcup_{k=1}^n V_k$ .

A closed subset of a compact set is compact.

**A.2.2 Images of compact sets under continuous functions are compact.**

**A.2.3 Uniform continuity.** Let  $(X, d)$  and  $(Y, \rho)$  be metric spaces and  $f$  a function from  $X$  to  $Y$ . Then  $f$  is called *uniformly continuous* if the following statement holds:

$$\forall \varepsilon > 0 : \exists \delta > 0 : \forall x, x' \in X : d(x, x') < \delta \Rightarrow \rho(f(x), f(x')) < \varepsilon.$$

If  $X$  is compact and  $f : X \rightarrow Y$  is continuous, then it is uniformly continuous.

**A.2.4 Sequences in compact metric spaces have convergent subsequences.**

**A.2.5 The Heine<sup>1</sup>-Borel<sup>2</sup> theorem.** The Heine-Borel theorem states that a subset of  $\mathbb{R}^n$  is compact if and only if it is closed and bounded. It is a very important result since it makes it easy to check compactness which in turn is an essential ingredient for several important results.

**A.2.6 The Bolzano<sup>3</sup>-Weierstrass theorem.** Suppose  $X$  is a metric space. Every sequence in  $X$  has a convergent subsequence if and only if  $X$  is compact.

## A.3. Product spaces

**A.3.1 Product spaces.** Let  $(X, d)$  and  $(X', d')$  be metric spaces. The function

$$\rho : (X \times X') \times (X \times X') \rightarrow [0, \infty) : ((x, x'), (y, y')) \mapsto \max\{d(x, y), d'(x', y')\}$$

is a distance function in  $X \times X'$  turning it into a metric space.

**THEOREM.** The cartesian product of compact metric spaces is compact.

---

<sup>1</sup>Eduard Heine (1821 – 1881)

<sup>2</sup>Émile Borel (1871 – 1956)

<sup>3</sup>Bernard Bolzano (1781 – 1848)

## Glossary

**annulus:** In a metric space an annulus is a set lying between two concentric circles. 16

**binary operation:** A binary operation on a set  $A$  is a function from  $A \times A$  to  $A$ . It is customary to express a binary operation as  $a \star b$  (or with other symbols in place of  $\star$ ). 3

**circumference of a triangle:** Two sum of the edge lengths of the triangle. 14

**congruence of triangles:** Two triangles are congruent if their edge lengths coincide. 14

**dense:** A subset  $A$  of a metric space  $(X, d)$  is called dense in  $X$  if every point of  $X \setminus A$  is a limit point of  $A$ . 17

**diameter:** In a metric space  $(X, d)$  the diameter of a set  $A \subset X$  is defined to be  $\sup\{d(x, y) : x, y \in A\}$ . 14

**group:** A set in which an associative binary operation is defined so that an identity and inverses to any element exist is called a group. 22

**identity:** An identity (in a set  $A$  with a binary operation  $\star$ ) is an element  $e \in A$  such that  $e \star a = a \star e = a$  for all  $a \in A$ . 3

**inverse:** The inverse of an element  $a$  (in a set  $A$  with a binary operation  $\star$  and identity  $e$ ) is an element  $b \in A$  such that  $a \star b = b \star a = e$ . 3

**isolated point:** A point  $x$  in a metric space  $(X, d)$  is called an isolated point of a set  $S \subset X$  if there exists an open disk  $B$  about  $x$  such that  $B \cap S = \{x\}$ . 7

**limit point:** A point  $x$  in a metric space  $(X, d)$  is called a limit point of a set  $S \subset X$  if  $B(x, 1/n) \cap S \setminus \{x\} \neq \emptyset$  for all  $n \in \mathbb{N}$ . 7

**linear:** A map  $F$  is called linear if  $F(\alpha u + \beta v) = \alpha F(u) + \beta F(v)$  for all  $\alpha, \beta \in \mathbb{C}$  and all  $u, v$  in the domain of  $F$ . 9

**local maximum:** The value  $u(z_0)$  is called a local maximum of a function  $u : S \rightarrow \mathbb{R}$  if  $u(z_0) \geq u(z)$  for all  $z$  in some neighborhood  $V = B(z_0, r) \cap S$  of  $z_0$ . 19

**monic:** A polynomial is called monic if its leading coefficient is equal to 1. 21

**monomial:** A function of the form  $m : \mathbb{C}^k \rightarrow \mathbb{C} : (z_1, \dots, z_k) \mapsto az_1^{n_1} \dots z_k^{n_k}$  when  $a$  is a complex number and  $n_1, \dots, n_k$  are non-negative integers. The number  $a$  is called the coefficient of the monomial while  $z_1, \dots, z_k$  are called the variables. 34

**multiset:** Let  $A$  be a set. A multiset is a function  $A \rightarrow \mathbb{N}_0$  (indicating how often an element  $a \in A$  is contained in the multiset). 9

- periodic:** A function  $f$  defined on  $\mathbb{C}$  (or  $\mathbb{R}$ ) is called periodic if there exists a complex (or real) number  $a \neq 0$  such that  $f(z + a) = f(z)$  for all  $z \in \mathbb{C}$  (or  $z \in \mathbb{R}$ ). The number  $a$  is called a period of  $f$ . 12
- permutation:** Let  $A$  be a finite set. A permutation of  $A$  is a bijective map from  $A$  to itself. 9
- polynomial:** A finite sum of **monomials**. 8, 34
- punctured disk:** A set is called a punctured disk if it is a disk with the center removed. 17
- rearrangement:** The series  $\sum_{n=1}^{\infty} w_n$  is a rearrangement of the series  $\sum_{n=1}^{\infty} z_n$  if there is bijective sequence  $k : \mathbb{N} \rightarrow \mathbb{N}$  such that  $w_n = z_{k_n}$ . 6
- similarity of triangles:** Two triangles are similar if the ratios of their respective edge lengths coincide. 14
- singleton:** A singleton is a set containing precisely one element. 5, 31
- symmetric polynomial:** A **polynomial** is called symmetric if no permutation of its variables changes its values. 21
- unit circle:** The set of points in  $\mathbb{C}$  of modulus 1. 22
- unit disk:** The set of points in  $\mathbb{C}$  of modulus less than 1. 11, 22
- zero:**  $x$  is called a zero of a function  $f$  if  $f(x) = 0$ . 12, 15

## List of Symbols

$|z|$ : the absolute value or modulus of the complex number  $z$ . 4

$\mathbb{C}$ : the set of complex numbers. 3

$\mathbb{C}_\infty$ : the Riemann sphere. 22

$\bar{z}$ : the complex conjugate of the complex number  $z$ . 3

$B(x_0, r)$ : the open disk of radius  $r$  centered at  $x_0$ . 31

$\overline{B}(x_0, r)$ : the closed disk of radius  $r$  centered at  $x_0$ . 31

$\text{Im}(z)$ : the imaginary part of the complex number  $z$ . 3

$n_f(r)$ : the number of zeros of an entire function  $f$  in the disk  $B(0, r)$ . 28

$\gamma^*$ : the range of the path or contour  $\gamma$ . 8

$\text{Re}(z)$ : the real part of the complex number  $z$ . 3

$Z(f)$ : the set of zeros of  $f$ . 15





## Hints

- 1.2.3:** Show that  $(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$  when  $n \in \mathbb{N}$ . **4**
- 1.2.6:** Assume the convex set  $S$  is not connected and pick points  $x \in S \cap A$  and  $y \in S \cap B$ . The fact that  $\{t \in [0, 1] : (1-t)x + ty \in A\}$  has a supremum leads to a contradiction. **5**
- 1.3.4:** The analogous statement is known to hold for real series. **6**
- 2.2.2:** Use property (1). **8**
- 2.3.3:** Let  $z = \int_a^b f \neq 0$  and  $\alpha = |z|/z$ . Then consider  $\int_a^b \alpha f$  which is real. **9**
- 2.3.8:** Given a point  $x$  in  $S$ , let  $C(x)$  denote the set of all  $y \in S$  for which there is a connected contour in  $S$  with initial point  $x$  and end point  $y$ . If  $S$  is open, then so are  $C(x)$  and  $S \setminus C(x)$ . **10**
- 2.5.2:** In trying to establish the derivative of  $z \mapsto f(z)$  at the point  $w$  let  $r = \max\{|z - z_0|, |w - w_0|\}$  and use the identity  $a^n - b^n = (a - b) \sum_{k=0}^{n-1} a^k b^{n-k-1}$ . **11**
- 2.5.5:**  $\phi(t) - z = (\phi(t) - z_0)(1 - (z - z_0)/(\phi(t) - z_0))$ . **12**
- 3.2.3:** One needs to show that the diameter of a triangle is the length of its longest edge. For this it is useful to realize that  $|x_1 - x_2| = |(r_1 - r_2)(a - c) + (s_1 - s_2)(b - c)| = |(s_1 - s_2)(b - a) + (t_1 - t_2)(c - a)| = |(t_1 - t_2)(c - b) + (r_1 - r_2)(a - b)|$  when  $x_j = r_j a + s_j b + t_j c$  and that two of the numbers  $r_1 - r_2$ ,  $s_1 - s_2$ , and  $t_1 - t_2$  are of the same sign unless one of them is zero. **14**
- 3.2.4:** Assume first that  $z_0 \notin \Delta$ . Let  $\Delta_0 = \Delta$ . Given  $\Delta_n$  let  $\Delta_{n+1}$  be one of the four triangles constructed in **3.2.3**. Then  $\bigcap_{n=1}^{\infty} \Delta_n$  consists of one point only, say  $w$ . By **2.2.2** the function  $h$  defined by  $f(z) = f(w) + f'(w)(z - w) + h(z)(z - w)$  is continuous and vanishes at  $w$ . Hence, given  $\varepsilon > 0$  and using **3.2.2** and **2.3.4**, there is an  $n$  such that  $|\int_{\Gamma_n} f| \leq 3\varepsilon D 4^{-n}$  when  $D$  denotes the diameter of  $\Delta$ . By making, at each step, the right choice of triangle in the above recursion one can also show that  $|\int_{\Gamma_n} f| \geq 4^{-n} |\int_{\Gamma} f|$ . This settles the first case. If  $z_0$  is a vertex of  $\Delta$ , we repeat the construction above but choosing  $\Delta_n$  always to be the one which has  $z_0$  as a vertex so that  $\int_{\Gamma} f = \int_{\Gamma_n} f$ . Finally, if  $z_0 \in \Delta \setminus \{a, b, c\}$  construct three triangles in  $\Delta$  which have  $z_0$  as a vertex. **14**
- 3.3.5:** Using **2.5.4** express  $f^{(k)}(z_0)$  in terms of a Taylor coefficient, then compute that coefficient employing a combination of **3.3.1** and **2.5.5**. **14**
- 3.3.9:** Define the set  $A$  of limit points of  $Z(f)$  in  $\Omega$  and show that both  $A$  and  $\Omega \setminus A$  are open. **15**
- 3.4.1:** Use **A.3.1** and **A.2.3**. **15**
- 3.4.2:** Note that  $\varphi(u, z) - \varphi(u, z_0) = \int_{(z_0, z)} \psi(u, \cdot)$ . Let  $h(z_0) = \int_{\Gamma} \psi(\cdot, z_0)$  and consider  $\left| \frac{g(z) - g(z_0)}{z - z_0} - h(z_0) \right|$ . **15**
- 3.4.3:** Note that  $f(u) - f(z) = \int_{(z, u)} f'(v) dv$  and  $f(u) - f(z) - f'(z)(u - z) = \int_{(z, u)} \int_{(z, v)} f''(w) dw dv$ . **15**

- 3.4.4:** On  $\Omega$  define  $g(z) = \int_{\Gamma} \varphi(u, z) du$  with  $\varphi$  as in 3.4.3. On  $\Omega' = \{z \in \mathbb{C} : \text{Ind}_{\Gamma}(z) = 0\}$  define  $h(z) = \int_{\Gamma} f(u)/(u - z) du$ . Then create an entire function by showing that  $h = g$  on  $\Omega \cap \Omega'$ . This function must be zero. **16**
- 3.4.7:** Fix  $z \in \Omega$ . For  $j = 1, 2$  let  $\gamma_j(t) = a + r'_j \exp(it)$  for  $t \in [0, 2\pi]$  and suitably chosen  $r'_j$ . Then consider  $\int_{\gamma_2 \ominus \gamma_1} f$ . To find  $a_n$  for  $n \geq 0$  employ 2.5.5; for the others use a variant of its proof after noting that  $u - z = -(z - a)(1 - (u - a)/(z - a))$ . **16**
- 4.1.2:** Define  $h(z_0) = 0$  and  $h(z) = (z - z_0)f(z)$  for  $z \neq z_0$ . Then show that  $h$  is holomorphic near  $z_0$ . **17**
- 4.1.6:** Assume on the contrary that there is a  $a \in \mathbb{C}$  and  $\delta > 0$  such that  $B(a, \delta) \cap f(B') = \emptyset$ . Then consider the function  $g(z) = 1/(f(z) - a)$  on  $B'$ . **17**
- 4.2.2:** Let  $m\gamma = \bigoplus_{k=1}^m \gamma$  and  $\Gamma' = \bigoplus_{k=1}^n \text{Ind}_{\Gamma}(z_k)\gamma_k$  where  $\gamma_k(t) = z_k + r_k \exp(it)$  for  $t \in [0, 2\pi]$  for sufficiently small but positive  $r_k$ . Then  $\int_{\Gamma} f = \int_{\Gamma'} f$ . **18**
- 4.2.3:** For the first integral consider a contour made up from the interval  $[-R, R]$  and a semi-circle of radius  $R$ ; for the second one consider a rectangle with vertices  $\pm R$  and  $\pm R + \pi i$ ; and for the last integral let  $u = \exp(it)$  and note that  $2i \sin(t) = u - 1/u$ . **18**
- 4.2.5:** Let  $\gamma$  parameterize the circle and set  $h = f/g$ . Then  $\int_{\gamma} h'/h = \text{Ind}_{h \circ \gamma}(0) = 0$  since  $(h \circ \gamma)^* \cap (-\infty, 0] = \emptyset$ . **19**
- 4.2.6:** Suppose  $f(z_0) = w_0 \in f(\Omega')$ . Let  $F = f - w_0$  and  $G = f - w$ . Then  $|F(z_0 + r \exp(it))| \geq \delta > 0$  for some  $r > 0$  and all  $t \in [0, 2\pi]$ . Determine how close  $w$  has to be to  $w_0$  so that Rouché's theorem 4.2.5 applies to  $F$  and  $G$ . **19**
- 5.1.2:** Use either Rouché's theorem or the fact that  $1/p$  would be entire if  $p$  had no zeros. **21**
- 5.1.3:**  $\prod_{j=1}^n z_j - \prod_{j=1}^n w_j = \sum_{j=1}^n q_j(z_j - w_j)$  for appropriate polynomials  $q_j$ . **21**
- 5.1.4:** If  $f(z) = \sum_{n=-\infty}^{\infty} a_n z^n$ , we have  $2\pi r_k^n a_n = \int_0^{2\pi} f(\gamma_k(t)) \exp(-int) dt$  where  $\gamma_k(t) = r_k \exp(it)$ . Hence  $\pi(r_k^n a_n + r_k^{-n} \overline{a_{-n}}) = \int_0^{2\pi} \text{Re } f(\gamma_k(t)) \exp(-int) dt$ . Now use, when  $n \in \mathbb{N}$ , that  $a_{-n} = 0$  and  $\int_0^{2\pi} C r_k^N \exp(-int) dt = 0$ . **21**
- 5.4.6:** Let  $F$  be an antiderivative of  $f'/f$  and consider the derivative of  $f \exp(-F)$ . **25**
- 5.4.7:** First note that  $\int_{-\varepsilon}^{\varepsilon} \log |1 - \exp(it)| dt$  is negligible when  $\varepsilon$  is small. For the remaining part note that  $\log |1 - z| = \text{Re}(\log(1 - z))$  when  $\text{Re}(z) < 1$ . **25**

# Index

- $n$ -th root, 4
- absolute convergence
  - of a series, 5
  - of an infinite product, 27
- absolute value, 4
- analytic, 11
- analytic continuation, 15
- associative law of addition, 3
- associative law of multiplication, 3
- ball, 31
- bounded, 31
- branch
  - of the logarithm, 24
- branch of the power function, 25
- canonical product, 29
- Cauchy sequence, 5
- closed, 31
- closed contour, 9
- commutative law of addition, 3
- commutative law of multiplication, 3
- compact, 32
- complex conjugate, 3
- complex plane, 4
- connected, 5
  - simply, 16
- continuity, 32
  - at a point, 32
- contour, 9
  - closed, 9
  - connected, 10
- convergence
  - of a function, 7
  - of a sequence, 31
  - of a series, 5
  - pointwise, 10
  - uniform, 10
- convex, 4
- cosine function, 23
- degree, 21
- derivative, 7
- differentiable, 7
- distance, 4, 31
- distance function, 31
- divergence
  - of a sequence, 32
  - of a series, 5
- edge of a triangle, 13
- elementary factor, 27
- end point
  - of a contour, 10
  - of a path, 8
- entire, 8
- essential singularity, 17
- exponent of convergence, 29
- exponential function, 12
- finite subcover, 32
- function
  - exponential, 23
  - rational, 22
- holomorphic, 8
- imaginary number, 3
- imaginary part, 3
- infinite product, 27
- initial point
  - of a contour, 10
  - of a path, 8
- Laurent series, 16
- length
  - of a line segment, 4
  - of a smooth path, 8
- limit of a sequence, 32
- line segment, 4
- logarithm, 24
- Möbius transform, 22
- meromorphic, 17
- metric, 31
- metric space, 31
- modulus, 4
- multiplicity

- of a pole, 17
  - of a zero, 15
- negative of a number, 3
- north pole, 22
- open, 31
- open cover, 32
- opposite of a path, 8
- order
  - of a pole, 17
  - of a zero, 15
- order of an entire function, 28
- partial sums, 5
- path, 8
  - smooth, 8
- polar coordinates, 4
- pole, 17
- polynomial, 21
- power, 4
- power series, 11
- primitive, 10
- principal part, 18
- radius of convergence, 11
- real part, 3
- reciprocal, 3
- residue, 18
- Riemann sphere, 22
- sequence
  - in a metric space, 31
  - of functions, 10
- sequence of partial sums, 5
- series, 5
  - geometric, 6
- simple pole, 18
- simply connected, 16
- sine function, 23
- singular part, 18
- singularity
  - essential, 17
  - isolated, 17
  - removable, 17
- sphere
  - unit, 22
- stereographic projection, 22
- topological space, 31
- topology, 31
- triangle, 13
- triangle inequality
  - for a metric, 31
  - for the absolute value, 4
- unit sphere, 22
- winding number, 13