

COMMUNITY
COORDINATED
MODELING
CENTER



NASA
Space
Weather
Research
Center


Flares, CMEs and SEPs

A. Taktakishvili

CCMC, SWRC

NASA Goddard Space Flight Center

In this presentation I will be talking about solar flares and coronal mass ejections and their relation to the space weather



Solar flare is a sudden brightening observed over the Sun's surface or the solar limb, which is interpreted as a large energy release. Flares are mainly followed by a mass ejections from the solar atmosphere called coronal mass ejections. The flare ejects clouds of electrons, ions, and atoms through the corona of the sun into space.


In this slide is shown the solar flare of 2012 July 12.

The three color images show SDO satellite (<http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Glossary>) images of the same flare in three different wave length intervals: 1) the left is SDO EVE X-ray image (http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Full_iSWA_Cygnets_List

The central panel shows SDO AIA extreme ultraviolet image for 13.1 nm http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/SDO_AIA_193 and the right panel - for a shorter wave length of 19.3 nm

http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/SDO_AIA_131

In this slide is shown also in the lower left corner a magnetogram (<http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Glossary>) - magnetic field map - of the solar photosphere for the same period of time. This flare is associated to the large and magnetically complicated active region in the southern hemisphere of the solar surface as is usually takes place.



In this slide is shown the time line for the X-ray flux intensity

(http://iswa3.cmc.gsfc.nasa.gov/wiki/index.php/GOES_X-RAY) for the same flare of 2012 July 12, in two different wave length intervals . A strong flare usually manifests itself by a sudden a jump of the X-ray intensity.

The flares are characterized by classes according to their X-ray flux intensity in the 0.1 – 0.8 nm wavelength range.

Flux/Wm² > 10⁻⁴ X - (eXtreme), strong

Flux/Wm² > 10⁻⁵ M – (Moderate)


Flux/Wm² > 10⁻⁶ C – (Common)

Flux/Wm² > 10⁻⁷ B – (Background)

Flux/Wm² > 10⁻⁸ A

M1.4 class flare that at the peak the flare intensity was equal to:

flux/Wm²=1.4*10⁻⁴




COMMUNITY
COORDINATED
MODELING
CENTER

Flares over the Solar cycle



Solar flares have been monitored by x-ray detectors on GOES satellites since 1976. The number of X-Class flares per month increases with the number of sunspots but **big flares can occur anytime sunspots are present.**



Solar flares have been monitored by x-ray detectors on GOES satellites since 1976.


In this plot the frequency of the X-class flares is plotted over the sunspot number plot. Green dots represent 1-2 X-class flares, yellow – 3-9 X-class flares and red – 10+ X class flares.

You can see that the number of X-Class flares per month increases with the number of sunspots, but big flares can occur anytime sunspots are present.



COMMUNITY
COORDINATED
MODELING
CENTER

Flare: SWx impacts




- Cause radio blackout through changing the structures/composition of the ionosphere (sudden ionospheric disturbances) – X ray and EUV emissions, **lasting minutes to hours**
- Affect radio communications, GPS, directly by its radio noises at different wavelengths
- Contribute to SEP – proton radiation, **lasting a couple of days**

5

Cause radio blackout through changing the structures/composition of the ionosphere (sudden ionospheric disturbances) – x ray and EUV emissions, **lasting minutes to hours**


Affect radio communications, GPS, directly by its radio noises at different wavelengths

Contribute to SEP – proton radiation, **lasting a couple of days**



COMMUNITY
COORDINATED
MODELING
CENTER

Flare Characteristics



- Flares tend to occur in isolation, localized in space and time but with strong correlations; typically one active region will produce dozens of flares, especially during periods of flux emergence (often near the beginning of the lifetime of a given region, but not always). **The most powerful events usually occur in active regions.**

Flares tend to occur in isolation, localized in space and time but with strong correlations; typically one active region will produce dozens of flares, especially during periods of flux emergence (often near the beginning of the lifetime of a given region, but not always). **The most powerful events usually occur in active regions.**




COMMUNITY
COORDINATED
MODELING
CENTER

Solar Flare Discovered



In 1859 Richard Carrington reported observing a large sunspot group on the afternoon of September 1st when "...two patches of intensely bright and white light broke out..."



In 1859 Richard Carrington reported observing a large sunspot group on the afternoon of September 1st when "...two patches of intensely bright and white light broke out...". The flare was followed by a major coronal mass ejection (CME) to travel directly toward Earth, taking 17.6 hours. This second CME moved so quickly because the first one had cleared the way of the ambient solar wind plasma.

On September 1–2, 1859, the largest recorded geomagnetic storm occurred. Aurorae were seen around the world, even over the Caribbean; those over the Rocky Mountains were so bright that their glow awoke gold miners, who began preparing breakfast because they thought it was morning. People who happened to be awake in the northeastern US could read a newspaper by the aurora's light.

Telegraph systems all over Europe and North America failed, in some cases shocking telegraph operators. Some telegraph systems continued to send and receive messages despite having been disconnected from their power supplies.




COMMUNITY
COORDINATED
MODELING
CENTER

Coronal Mass Ejection



Coronal Mass Ejection – Reaches the Earth in 1-3 days



From time to time huge mass of solar plasma can erupt from the solar atmosphere. This phenomenon is called coronal mass ejection (CME,

<http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Glossary>).

If CME is directed towards Earth, depending on it's speed and size, it can reach the Earth in 1-3 days. CMEs are usually causing the strongest geomagnetic storms.



COMMUNITY
COORDINATED
MODELING
CENTER


CME/flare



- The most energetic CMEs occur in close association with powerful flares. Nevertheless large-scale CMEs do occur in the absence of major flares even though these tend to be slower and less energetic.
- When strong flare/CME occurs, it gives off emission across the whole electromagnetic spectrum, at the same time energetic particles


The most energetic CMEs occur in close association with powerful flares. Nevertheless large-scale CMEs do occur in the absence of major flares even though these tend to be slower and less energetic.

When strong flare/CME occurs, it gives off emission across the whole electromagnetic spectrum, at the same time energetic particles



In this slide is shown the solar flare of 2012 July 12 yet again and the following CME in the SDO EUV 193 image.


You can see the motion of the material and opening of the coronal loops in the area adjacent to the flare and the active region.



But the better view of a CME and its motion in the interplanetary space can be obtained from the coronagraph images. Coronagraphs create an artificial eclipse of the sun. Eclipses allow corona to be better viewed, but natural eclipses do not happen often. Occulting disk blocks the bright sun so we can observe corona features.

This slide shows the same CME seen in 3 coronagraphs located on three different satellites. The schematic in the upper center shows the location of STEREO B, SOHO and STEREO A satellites in the ecliptic plane at the time when the CME occurred.

The CME was moving sort of towards the Earth, so STB sees it moving to the right in its plane of sky, for STA it moves to the left and it's a halo image for SOHO that is located on a sun-earth line.



Not all CMEs originate from the active regions. Here is shown an example of a CME caused by so called filament eruption. Filaments are formed in magnetic loops that hold relatively cool, dense gas suspended above the surface of the Sun. Magnetic instabilities cause filaments to raise and parts of it disconnect from the solar surface resulting into CMEs.



COMMUNITY
COORDINATED
MODELING
CENTER

CME Properties



- Mass: $\sim 10^{15-16}$ g
- Speed: few hundred - 3000 km/s

..or

- Mass: ~ 1 million Nimitz-class aircraft carriers
- Speed: 1.5 - 10 million km/hour



Earth?



- Arrives to Earth in 1-2 days

CME properties:

- Mass: $\sim 10^{14}$ kg
- Speed: few hundred - 3000 km/s

In other words:

- Mass: ~ 1 million Nimitz-class aircraft carriers
- Speed: 1.5 - 10 million km/hour

If it hits the Earth, it can have a significant impact.




COMMUNITY
COORDINATED
MODELING
CENTER

CME SCORE



- A simple new category system for CMEs based on frequency of detection and speed
- Complements Flare Classes
- Applicable in space weather operations and research

Space Weather Research Center CME SCORE Scale



Recently we, here at SWRC, introduced, as we call it, CME SCORE scale, a simple new category system for CMEs based on frequency of detection and speed.

It Complements Flare Classes and is applicable in space weather operations and research.

The graph here shows the frequency of the CME detection (number of events per year) as a function of the CME speed. Bin size is 100 km/s.

So we have S-types CMEs with the speed less than 500/km/s, C-type (common) for the range 500-999, O-type (occasional) for 1000-1999, R-type (rare) for 2000-2999 and ER-type (extremely rare) CMEs for CMEs with the speed more than 3000 km/s.



COMMUNITY
COORDINATED
MODELING
CENTER

SWx impacts of CME



- Contribute to SEP (particle radiation): 20-30 minutes from the occurrence of the CME/flare
- Result in a geomagnetic storm: takes 1-2 days arriving at Earth
- Result in electron radiation enhancement in the near-Earth space: takes 1-3 days

Affecting spacecraft electronics – surfacing charging/
internal charging, radio communication, navigation
power grid, pipelines, and so on

15

Contribute to SEP (particle radiation): 20-30 minutes from the occurrence of the CME/flare


Result in a geomagnetic storm: takes 1-2 days arriving at Earth

Result in electron radiation enhancement in the near-Earth space: takes 1-3 days.


Contribute to SEP (particle radiation): 20-30 minutes from the occurrence of the CME/flare

Result in a geomagnetic storm: takes 1-2 days arriving at Earth

Result in electron radiation enhancement in the near-Earth space: takes 1-3 days



When the CME arrived to the ACE satellite, close to the Earth, it caused a shock seen here in the magnetic field and the solar wind speed measurements by ACE. This impact caused a strong disturbance in the Earth's magnetosphere or as they call it geomagnetic storm.



The strength of the geomagnetic storm is measured by so called Kp index, which characterizes degree of the disturbance of the Earth' magnetic field ground base measurements. Kp goes from 0 to 9. This storm with Kp=6 is considered to be a moderate one.




COMMUNITY
COORDINATED
MODELING
CENTER


Physical Mechanism Behind the Flares and CMEs



It is believed that solar magnetic field, releases energy, accelerating solar plasma and causing flares and CMEs. Magnetic reconnection.



What is the mechanism behind the flares and CMEs. It is believed that the magnetic field can change its configuration in a constantly varying solar atmosphere and during this reconfiguration it releases energy accelerating solar plasma causing flares and CMEs. The scientists are still debating on the details of the mechanisms, but the fact that the magnetic field is involved somehow is accepted by everybody.



This movie shows ENLIL modeling of the same July 12, 2012 CME. In all three panels normalized solar wind density is plotted.

In the left panel is shown view from the Solar north pole down to the ecliptic plane, where all the planets are moving around the sun. Yellow circle is the Earth, orange (closest to the Sun) – Mercury, green – Venus, red – Mars, red and blue squares – STEREO A and B satellites respectively.

The second panel is a meridional cut passing through the Earth trajectory. The latitude goes from > -60 to $< +60$ degrees.

The third panel shows Latitude–Longitude map of a sphere at 1 AU, cut for latitudes < -60 and $> +60$ degrees.

You can see that the CME causes enormous disturbance throughout the whole heliosphere. CMEs are called sometimes Space Hurricanes. For the terrestrial (regular) weather we need to know the possible path of a hurricane and its strength. The similar way, for the Space Weather we need to know the CME propagation path and its strength (speed, mass content) to estimate its possible impact.



COMMUNITY
COORDINATED
MODELING
CENTER

Solar Energetic particles - What are they?



Definition:

Energetic charged particles (such as electrons and protons) traveling much faster than ambient particles in the space plasma, at a % of the speed of light (relativistic!)

Elemental composition* (may vary event by event)

96.4 % protons

3.5% alpha particles

0.1% heavier ions (not to be neglected!)

Energies: up to ~ GeV/nucleon

They can travel from the Sun to the Earth in one hour or less!

The term SEP usually refers to protons (even though “p” is particle)

An SEP event is an enhancement in the radiation environment.

Summarize effects again.


Elemental composition* (may vary event by event)

96.4% protons

3.5% alpha particles

0.1% heavier ions (not to be neglected!)


Energies: up to ~ GeV/nucleon



Particles are accelerated during the flares and CMEs. When fast CME moves in the heliosphere it creates a shock. In front of the shock particles are accelerated resulting into SEP events observed at Earth and satellites. Not all the observed flares and CMEs lead to SEP observation. The flare and CME front locations have to be magnetically connected to the observation point.

When flare and CME occurs accelerated charged particles start to move along the magnetic field lines. The field lines are like the railroad for a train, where charged particles depart from the solar atmosphere and whether

they arrive to the Earth or STEREO B, for example, depends on the fact whether if the interplanetary magnetic field they are moving along is connected to the Earth/STEREO B or not.



COMMUNITY
COORDINATED
MODELING
CENTER

How do we monitor SEP Levels



Track the particle flux at different locations.

Units: pfu, pfu/MeV

(1 pfu = 1 particle flux unit = $1/\text{cm}^2/\text{sec}/\text{sr}$)

- **STEREO In-situ Measurements of Particles and CME Transients (IMPACT)**
 - *Differential energy band; example energy range: 13-100 MeV*
- **Upstream of Earth with SOHO/COSTEP**
 - *Differential energy bands; example energy range: 15.8-39.8 MeV*
- **Geostationary Orbit with GOES**
 - *Integral flux, example energy ranges: >10 MeV, >100 MeV*


Fluence = flux integrated over the entire event - dose

Important for biological effects (flights)

Event magnitudes:

- > 10 MeV/nucleon integral fluence: can exceed 10^9 cm^{-2}
- > 10 MeV/nucleon peak flux: can exceed $10^5 \text{ cm}^{-2}\text{s}^{-1}$

Converting between the two is important to know in order to quantify the relative strengths.



Another way that we “see” SEPs are in images, for example, coronagraphs. Here you can see the ‘snow’ of particles hitting the detector.



COMMUNITY
COORDINATED
MODELING
CENTER

How do we define an SEP Event?




At the SWRC, SEP events are defined as:

GOES Proton $E > 10$ MeV channel > 10 pfu

GOES Proton $E > 100$ MeV channel > 1 pfu


SOHO Proton, > 15.8 MeV channel > 0.1 pfu/MeV

STEREO Impact 13-100 MeV channel > 0.1 pfu/MeV



Here is an example of an event where the 10 MeV limit was met, but not the 100 MeV.

SOHO: > 15.8 MeV proton channels 0.1 pfu/MeV.



If you are speaking about one specific location in the heliosphere, for example, just at the Earth, then during solar minimum, there can be only a few or even none in a year. During higher activity times, there can be about one per month or so.

However, there can be many more SEP events happening at other locations, or that we simply are not measuring – remember how that sometimes events can be focused in space!

With only four events in the first 5 months of 2013, things are a little more quiet than we would have guessed.






COMMUNITY
COORDINATED
MODELING
CENTER

Can we forecast SEP events?




Uses detection of high energy *electrons* to predict arrival of high energy *protons*

Data source: SOHO/
COSTEP




http://iswa.gsfc.nasa.gov/wiki/index.php/SOHO/Costep_Proton_Flux_-_Forecast




COMMUNITY
COORDINATED
MODELING
CENTER


Profile Shapes of SEP Fluxes



Impulsive SEP event - The peak at the beginning due to flare, all off – indicates how well connected you are to the source (timing)

Gradual SEP event - Slow rise, then peak when the ICME passes the spacecraft






Not only were the >10 Mev channel elevated well above our threshold, but the >50 and >100 channels as well, which tell us that this event was very efficient at accelerating the highest energy particles.

So, we always watch for Active Regions passing on the Western limb, behind the Sun.

This event shown here, May 17, was very very strong...lead into GLE slide



The most extreme events...

This suspicion was confirmed by the detection of neutron enhancement at 4 stations. Those neutrons are reaction particles from relativistic protons

Add arrows to mark stations that observed May 17 GLE – what WAS THE TIME????

Save discussion of GLE events in SC 23 and characteristics of GLEs until 2nd half of talk. First half should only be observations!



What causes strong SEP events?



Complexity of AR

- Most young, more compact
- Magnetic connectivity of AR
 - About ~50% are well connected
- Magnitude of flare
 - Average X3.8, but as low as M7.1
 - Long duration
- Magnitude of CME
 - Range of speeds (~2,000 km/s average, but four events <1,500 km/s)
- Seed particles
 - Known to have harder spectrum

Table 1. GLE events and associated flares and CMEs (adopted from Gopalswamy et al. 2010)

GLE ID	Date	Time ^a	Max Int (%) ^b	Flare		CME POS speed (km/s)	Width (deg)
				GOES Class	Location		
55	1997/11/06	12:10	11.3	X9.4	S18W63	1556	360
56	1998/05/02	13:55	6.8	X1.1	S15W15	938	360
57	1998/05/06	08:25	4.2	X2.7	S11W66	1099	190
58	1998/08/24	22:30	3.3	X1.0	N35E09	3 ^c	-3 ^c
59	2000/07/14	10:30	29.3	X5.7	N22W07	1674	360
60	2001/04/15	14:00	56.7	X14	S20W85	1199	167
61	2001/04/18	02:35	18.8	C2.2	S20W116	2465	360
62	2001/11/04	17:00	3.3	X1.0	N06W18	1810	360
63	2001/12/26	05:30	7.2	M7.1	N05W54	1446	>212
64	2002/08/24	01:18	5.1	X3.1	S02W81	1913	360
65	2003/10/28	11:22	12.4	X17	S18E18	2459	360
66	2003/10/29	21:30	8.1	X10	S18W04	2029	360
67	2003/11/02	17:30	7.0	X8.3	S18W57	2598	360
68	2005/01/17	09:55	3.0	X3.8	N14W25	2547	360
69	2005/01/20	06:51	277.3	X7.1	N14W61	3242 ^c	360
70	2005/12/13	02:45	92.3	X3.4	S06W23	1774	360

^aAccording to the Oulu Neutron Monitor
^bNo SOHO LASCO data
^cFrom Gopalswamy et al. (2010). There are different estimates (see Grechnev et al. 2008)

Nitta et al. 2012

Gopalswamy et al. 2012, Li et al. 2012, Mewaldt et al. 2012

Like I said, these events prompt more questions and more questions, so we want to look at all of the extreme events (GLE) and see if any characteristics of the solar part of the storm can tell us about what is happening. And the answer is that while there are trends, it's not true that strong flare cause GLEs, as you saw only M5 cause the May event, it's not the fastest and biggest CMEs.

Again I want to emphasize that these are all just speculations because most of the interesting stuff is happening in a region that is not well covered by coronagraphs.

End with: And so we are motivated to model the acceleration of particles by CME driven shocks in the low corona...

Comparing to previous AR1476 flares: Longer duration, but similar magnitude

March 7th event: flare and CME weaker, SEPs at Earth 20x weaker

Solar Cycle 23 GLE events: no CME in previous 24 hours (to generate seed particles)

What was special about this flare and CME to generate a GLE?

Timing: CME erupted during rise time of flare


Connectivity: AR well connected to Earth

CME speed 1,500 km/s sufficient to drive shock in low corona (Evans et al. 2008)


Must have been very wide CME (to impact STEREO A and Earth)

As is common, they were not geoeffective (Kp max =4)

Possible Type II radio burst indicates CME driven shock



Homework



- In SDO AIA 193 image find the location of the flare that started around 2013-03-15T06:00
- What was the class of the flare?
- Was the flare accompanied by a CME?
- If yes, was the CME Earth directed?
- What was the start time of the CME in different coronagraph images?
- Was there an SEP event observed related to the flare and CME?
- Did the CME arrive to the Earth (ACE)?
- If yes, when did it arrive?
- What was the Kp geomagnetic storm index due to the CME arrival?
- Was this CME modeled?
- When was the predicted arrival of the CME?