

# A GENERALIZATION OF A THEOREM OF CARNOT

Florentin Smarandache  
 University of New Mexico  
 200 College Road  
 Gallup, NM 87301, USA  
 E-mail: smarand@unm.edu

**Theorem of Carnot:** Let  $M$  be a point on the diagonal  $AC$  of an arbitrary quadrilateral  $ABCD$ . Through  $M$  one draws a line which intersects  $AB$  in  $\alpha$  and  $BC$  in  $\beta$ . Let us draw another line, which intersects  $CD$  in  $\gamma$  and  $AD$  in  $\delta$ . Then one has:

$$\frac{A\alpha}{B\alpha} \cdot \frac{B\beta}{C\beta} \cdot \frac{C\gamma}{D\gamma} \cdot \frac{D\delta}{A\delta} = 1.$$

**Generalization:** Let  $A_1 \dots A_n$  be a polygon. On a diagonal  $A_1 A_k$  of this polygon one takes a point  $M$  through which one draws a line  $d_1$  which intersects the lines  $A_1 A_2, A_2 A_3, \dots, A_{k-1} A_k$  respectively in the points  $P_1, P_2, \dots, P_{k-1}$  and another line  $d_2$  intersects the other lines  $A_k A_{k+1}, \dots, A_{n-1} A_n, A_n A_1$  respectively in the points  $P_k, \dots, P_{n-1}, P_n$ . Then one has:

$$\prod_{i=1}^n \frac{A_i P_i}{A_{\varphi(i)} P_i} = 1,$$

where  $\varphi$  is the circular permutation

$$\begin{pmatrix} 1 & 2 & \dots & n-1 & n \\ 2 & 3 & \dots & n & 1 \end{pmatrix}.$$

*Proof:*

Let us have  $1 \leq j \leq k-1$ . One easily shows that:

$$\frac{A_j P_j}{A_{j+1} P_j} = \frac{D(A_j, d_1)}{D(A_{j+1}, d_1)}$$

where  $D(A, d)$  represents the distance from the point  $A$  to the line  $d$ , since the triangles  $P_j A_j A'_j$  and  $P_j A_{j+1} A'_{j+1}$  are similar. (One notes with  $A'_j$  and  $A'_{j+1}$  the projections of the points  $A_j$  and  $A_{j+1}$  on the line  $d_1$ ).

It results from it that:

$$\frac{A_1 P_1}{A_2 P_1} \cdot \frac{A_2 P_2}{A_3 P_2} \dots \frac{A_{k-1} P_{k-1}}{A_k P_{k-1}} = \frac{D(A_1, d_1)}{D(A_2, d_1)} \cdot \frac{D(A_2, d_1)}{D(A_3, d_1)} \dots \frac{D(A_{k-1}, d_1)}{D(A_k, d_1)} = \frac{D(A_1, d_1)}{D(A_k, d_1)}$$

In a similar way, for  $k \leq h \leq n$  one has:

$$\frac{A_h P_h}{A_{\varphi(h)} P_h} = \frac{D(A_h, d_2)}{D(A_{\varphi(h)}, d_2)}$$

and

$$\prod_{h=k}^n \frac{A_h P_h}{A_{\varphi(h)} P_h} = \frac{D(A_k, d_2)}{D(A_1, d_2)}$$

The product of the theorem is equal to:

$$\frac{D(A_1, d_1)}{D(A_k, d_1)} \cdot \frac{D(A_k, d_2)}{D(A_1, d_2)},$$

but

$$\frac{D(A_1, d_1)}{D(A_k, d_1)} = \frac{A_1 M}{A_k M}$$

since the triangles  $MA_1A_1'$  and  $MA_kA_k'$  are similar. In the same way, because the triangles  $MA_1A_1''$  and  $MA_kA_k''$  are similar (one notes with  $A_1''$  and  $A_k''$  the respective projections of  $A_1$  and  $A_k$  on the line  $d_2$ ), one has:

$$\frac{D(A_k, d_2)}{D(A_1, d_2)} = \frac{A_k M}{A_1 M}.$$

The product from the statement is therefore equal to 1.

Remark: If one replaces  $n$  by 4 in this theorem, one finds the theorem of Carnot.